

## First report of leech predation on *Pleurodeles nebulosus* (Guichenot, 1850) in Kabylia, Algeria

Khaled Merabet, Abdelhak Dahmana, Mokrane Karar & Aissa Moali

Laboratoire de recherche en Ecologie et Environnement, Faculté des Sciences de la Nature et de la Vie, Université de Bejaia, 06000 Bejaia, Algeria. C.e. : kaled.merabet@gmail.com

Fecha de aceptación: 8 de febrero de 2017.

Key words: Kabylia, Algerian ribbed newt.

**RESUMEN:** En la presente nota se describe un caso de parasitismo del hirudíneo *Hirudo troctina* sobre el urodelo *Pleurodeles nebulosus* en Kabylia.

In Algeria, the Algerian newt, *Pleurodeles nebulosus* (Guichenot, 1850), is present in humid, sub-humid and semi-arid Mediterranean areas but absent from the northwestern region (Mateo *et al.*, 2013). The distribution map for *P. nebulosus* largely follows the map presented in Veith *et al.* (2004). Recently, new studies provided some valuable data on the ecology and reproduction of *P. nebulosus* in Algeria (e.g., Ben Hassine & Escoriza, 2014; Escoriza & Ben Hassine, 2015; Merabet *et al.*, 2016). The continuing decline in the extent and quality of amphibians' habitat are the main cause of amphibian populations decline in the Maghreb (e.g., Samraoui *et al.*, 2012). Between other minor threats, intensive leech predation has been reported as a potential cause for local population decline (Beukema & Philip de Pous, 2011).



**Figure 1:** a) Dead individual of *P. nebulosus* in Chemini. b) The visited site in Chemini (Kabylia). c) Predation by *H. troctina* on *P. nebulosus* in Chemini (Kabylia).

**Figura 1:** a) Ejemplar de *P. nebulosus* muerto en Chemini. b) El lugar visitado en Chemini (Kabylia). c) Depredación de *P. nebulosus* por *H. troctina* en Chemini (Kabylia).

Predation of leeches upon *P. nebulosus* is a known and already reported phenomenon from Tunisia (Ben Hassine *et al.*, 2013). However, is not yet well understood if this predation is a common phenomenon and can have a negative effect on amphibian populations. In Algeria, based on the existing literature, Billet (1904) was the first to mention leech predation on populations of *Pelophylax saharicus* (Boulenger, 1913), who related the presence of the leech with anuran trypanosomiasis.

With this note, we wish to draw attention to one particular parasite, *Hirudo troctina* (Johnson, 1816) which may be a locally important source of mortality for *P. nebulosus* in Kabylia.

While conducting a survey in Kabylia (Chemini) on 14 March 2016, we made several observations of dead individuals of

*P. nebulosus* (Figure 1a) near a large permanent pound situated at 1600 masl ( $36^{\circ}37'41.406''$  N,  $4^{\circ}34'4.705''$  E) (Figure 1b). In the visited site we found 10 dead animals and many others suffering from leech predation by *H. troctina* (Figure 1c). After a thorough exploration of the water body we found a lot of leeches attacking especially *P. nebulosus* and no other amphibians.

In conclusion, we propose that punctual predation episodes by freshwater leeches (*H. troctina*) in the reported site can cause many deaths in *P. nebulosus* populations at the local scale. This observation should be taken seriously to measure if these attacks by leeches have effects on local population dynamics, and a strict study is needed to know if it is affecting other *P. nebulosus* populations.

## REFERENCES

- Ben Hassine, J. & Escoriza, D. 2014. New ecological data on the family Salamandridae in the Maghreb. *Herpetological Review*, 45:193–200.
- Ben Hassine, J., Kassabi, A. & Nouira, S. 2013. *Pleurodeles nebulosus* (Guichenot, 1850) en Tunisie: répartition, habitat, reproduction et statut. *Bulletin de la Société Herpétologique de France*, 144: 51-66.
- Beukema, W. & Pous, P. 2010. Exceptional leech predation on *Amietophryne mauritanicus* (Anura, Bufonidae) in Tunisia. *Herpetology Notes*, 3: 289-290.
- Billet, A. 1904. Culture d'un Trypanosome de la grenouille chez une hirudine; relation ontogénique possible de ce Trypanosome avec un Hemogregarine. *Comptes rendus de l'Académie des Sciences*, 137: 574-76.
- Escoriza, D. & Ben Hassine, J. 2015. Niche Partitioning at Local and Regional Scale in the North African Salamandridae. *Journal of Herpetology*, 49: 276-283.
- Mateo, J.A., Geniez, P. & Pether, J. 2013. Diversity and conservation of Algerian amphibian assemblages. 51-84. In: Busack, S. & Heatwole, H. (eds.), *Amphibian Biology. Volume 11. Status of Conservation and Decline of Amphibians: Eastern Hemisphere Part 2*. Asociación Herpetológica Española. Madrid.
- Merabet, K., Dahmana, A., Karar, M. & Moali, A. 2016. New occurrence record of the Algerian ribbed newt *Pleurodeles nebulosus* (Guichenot, 1850) in Algeria. *Herpetological Bulletin*, 137: 43.
- Samraoui, B., Samraoui, F., Benslimane, N., Alfarhan, A. & Al-Rasheid, K.A.S. 2012. A precipitous decline of the Algerian newt *Pleurodeles poireti* (Gervais, 1835) and other changes in the status of amphibians of Numidia, north-eastern Algeria. *Revue d'Écologie (Terre & Vie)*, 67: 71-81.
- Weith, M., Mayer, C., Samraoui, B., Donaire Barroso, D. & Bogaerts, S. 2004. From Europe to Africa and vice versa: evidence for multiple intercontinental dispersal in ribbed salamanders (genus *Pleurodeles*). *Journal of Biogeography*, 31: 159-171.