

Ten years of research on Bolivian amphibians: updated checklist, distribution, taxonomic problems, literature and iconography

IGNACIO DE LA RIVA¹, JÖRN KÖHLER², STEFAN LÖTTERS² & STEFFEN REICHLE²

¹ Museo Nacional de Ciencias Naturales, CSIC,

C/ José Gutiérrez Abascal 2, 28006 Madrid, Spain

² Zoologisches Forschungsinstitut und Museum Alexander Koenig,
Adenauerallee 160, 53113 Bonn, Germany.

Abstract: After ten years of intensive research, an updated checklist of the amphibians of Bolivia was compiled. For each species, information on its general distribution within the Neotropics, its occurrence within Bolivian departments and ecoregions, and habitat use is provided. Key literature and taxonomic remarks, if pertinent, are added. The list contains 186 valid species recorded from Bolivian territory, 33 (17.7%) of which are Bolivian endemisms. Two species reported recently (*Hyla walfordi* and *Phrynobatus peruvianus*) are not included because those reports were based on wrong identifications. Nine species (*Allobates femoralis*, *Epipedobates trivittatus*, *Hyla koechlini*, *H. sarayacuensis*, *Phrynohyas resinifictrix*, *Scinax squalirostris*, *Phyllomedusa tomopterna*, *Eleutherodactylus ventrimarmoratus*, and *Ischnocnema quixensis*) are recorded for the first time. Moreover, 67 additional species are expected to occur in the country. Finally, we provide a comprehensive bibliography on Bolivian amphibians and an iconography, figuring almost all Bolivian species in color.

Keywords: Amphibia, Bolivia, checklist, taxonomy, distribution, bibliography, iconography

Resumen: Diez años de investigación sobre anfibios de Bolivia: lista actualizada, distribución, problemas taxonómicos, bibliografía e iconografía.- Se presenta una lista actualizada de los anfibios de Bolivia, que recopila los resultados de diez años de investigación. Para cada especie se proporciona información sobre distribución general en el Neotrópico y en los departamentos y ecoregiones bolivianas, uso del hábitat y bibliografía clave, así como comentarios taxonómicos, cuando proceden. La lista contiene 186 especies válidas registradas en territorio boliviano, de las cuales 33 (17.7%) son endémicas. Dos especies citadas recientemente (*Hyla walfordi* y *Phrynobatus peruvianus*) no se incluyen en la lista porque dichas citas se basaron en identificaciones erróneas. Nueve especies (*Allobates femoralis*, *Epipedobates trivittatus*, *Hyla koechlini*, *H. sarayacuensis*, *Phrynohyas resinifictrix*, *Scinax squalirostris*, *Phyllomedusa tomopterna*, *Eleutherodactylus ventrimarmoratus*, e *Ischnocnema quixensis*) se citan en este trabajo por primera vez. Además, 67 especies aún no citadas podrían existir en el país. Finalmente, se proporciona una bibliografía exhaustiva sobre los anfibios de Bolivia y fotografías en color de casi todas las especies registradas.

Palabras clave: Amphibia, Bolivia, lista taxonómica, taxonomía, distribución, bibliografía, iconografía

INTRODUCTION

For many decades, the amphibian fauna of Bolivia has been one of the least known in the world. Until 1990, the information available was scattered among several publications devoted mainly to original descriptions of species, reports on collecting trips, and revisions of several taxa which included Bolivian material. The first comprehensive list of Bolivian amphibians was provided by DE LA RIVA (1990a), who compiled most of the published information, arranged the available data on

distribution, included data from several collections, added some new findings, and provided color photographs of 62 species, some of them never illustrated before. This list contained 112 species accepted as valid, plus several others of doubtful taxonomic status. In addition, a total of 47 species from four different biogeographic areas were considered as likely occurring in the country.

Throughout the 1990s, the Bolivian amphibians have been the subject of intensive research, which is obvious from the increased number of recent publications (see biblio-

graphic section below). As a result, the number of species known for the country by early 2000 is much higher than the 1990 number. This fact means not only a dramatic improvement of the knowledge of the diversity of Bolivian amphibians, but also of that of their taxonomy, distribution, ecology, and natural history. Especially remarkable are the many publications dealing with vocalizations, which have led to the description of the calls of nearly 100 Bolivian species of anurans.

We are convinced of the importance of papers summarizing the state of the art of the distribution and diversity of regional faunas, especially those of poorly surveyed countries. The first step is "to know what is known". Thus, these summaries, although best considered works in-progress, are essential not only as keystones on which the advance of knowledge, future research and conservation policies must be based, but also as an important stimulus and guidance for students and other people interested in natural resources. Following this philosophy and given the necessity of a new compilation of the huge amount of new information generated in the last few years, we have prepared the present contribution, whose main goals are to: (1) provide an updated checklist of Bolivian amphibians; (2) give brief guidelines about their geographic and ecological distributions; (3) point out particular cases in which taxonomic problems preclude accurate or definitive statements; (4) include a comprehensive bibliography; (5) discuss the present results and predict plausible, future findings; and (6) show color photographs of almost every amphibian species in Bolivia, many of them illustrated herein for the first time.

MATERIAL, METHODS AND GUIDELINES

Unless stated otherwise, the information provided in the list mainly comes from published papers cited in the literature, which has been updated until April 2000, and also in-

cludes papers in press about which we are aware. Basically we intend to reflect the current state of taxonomy of Bolivian amphibians. Thus, in general, in this paper we do not make taxonomic decisions or changes although we are aware of some cases in which such changes would be necessary and justified. Several species are reported for the first time in the present paper, based mostly on voucher specimens. Records based solely on personal communications by other persons have not been included, unless they have been corroborated by us. Apart from published information, additional data from collections and the personal experience of the authors have been used to generate the patterns of geographic and ecological distributions of the species. Orders, families, subfamilies (when pertinent), genera, and species are arranged in alphabetical order. Each species entry consists of the scientific name, author, and year of description, which is followed by an asterisk when the type locality is in Bolivia, and two asterisks when the species is endemic to the country. Additional information is arranged into three sections: "Distribution", "Key literature" and "Remarks".

Distribution.- Under this heading, for each species, the following five kinds of information are provided:

(1) Patterns of distribution in the Neotropics.- They are general geographic areas shared by several species with similar ecological requirements and/or biogeographical histories. Some broad patterns encompass other, more restricted ones. The patterns are:

Amazonia, which comprises the entire Amazon basin plus, usually, the Upper Orinoco basin and the Guianan region.

Andean Highlands, which includes the puna and Altiplano from, at its greatest extension, S Peru to N Argentina and Chile.

Bolivian-Argentinean Forests, comprising the semi-humid montane Andean forests from Santa Cruz (Bolivia) to N Catamarca (Argentina).

Cerrado domain, which comprises the open formations whose core area lies in Central and SE Brazil and may also include open areas in the lower Amazon and NE Brazil (Caatingas), reaching the Guianas, and/or the Plata Basin in Argentina and adjacent countries.

Chaco, which extends mainly throughout N Argentina, Paraguay, SW Brazil and SE Bolivia. The Chaco is often divided into dry and humid Chaco. The later holds some species which are widespread in other areas or are otherwise linked to tropical environments, whereas the dry Chaco is inhabited by the typical Chacoan species adapted to the extreme conditions of this ecosystem.

Humid Montane Forests, and/or humid subparamo in the Andes from S Peru to Bolivia or only those of Bolivia.

Lower Amazon Basin, which comprises the eastern half of the Amazon Basin and may include the Guianan region.

Middle and South America, which corresponds to species with a broad distribution that may reach up to Mexico in the north and SE Brazil and northern or central Argentina in the south, albeit more commonly these distributions range from Panama or Costa Rica to Bolivia, comprising both tropical and subtropical environments (often such broad distributions reflect a poor knowledge on the systematics of a given species complex and, as this knowledge improves, less species will fall within this pattern of distribution).

Peri-Andean Forests, which is the pattern of those species occurring in the foothills and lower Andean slopes from Colombia or Ecuador to Bolivia, but more often from SE Peru to Bolivia.

South America, which comprises tropical and subtropical regions from Colombia to SE Brazil and northern or central Argentina.

Southwestern Amazonia, comprising roughly the Madeira river basin in SE Peru, Amazonian Bolivia, and the Brazilian states of Acre and Rondônia.

Subtropical South America, comprising

temperate and subtropical areas south of the Chaco and the Cerrado, in Argentina, Bolivia, SE Brazil, Paraguay and Uruguay. Bolivian species with this general pattern usually occur only in the inter-Andean valleys.

Temperate Inter-Andean Valleys of Bolivia and adjacent areas of N Argentina.

Tropical Middle and South America, including Costa Rica and/or Panama, the Amazon and Orinoco basins and the Guianas, and may include the Brazilian Atlantic forest; it may also include the Pacific lowlands of Colombia and Ecuador.

Tropical South America, which is like the previous one but excluding Middle America.

Upper Amazon Basin, which comprises roughly the western half of the Amazon Basin and may include or not the Guianan region.

All these patterns represent only a rough way to make species distributions fit in a scheme. Consequently, some species can be assigned to more than one region or cannot be easily assigned to any of these regions.

(2) Bolivian departments where the species occurs. The nine Bolivian departments are (Figure 1): Beni, Cochabamba, Chuquisaca, La Paz, Oruro, Pando, Potosí, Santa Cruz, and Tarija. For each species, those departments for which we are aware of published records, voucher specimens or personal observation by any of the authors, are marked in boldface. If the presence of the species is not confirmed but only expected (for example, because it occurs in neighboring areas of other departments or the type of habitat is the same), the department appears in normal letters.

(3) Bolivian ecoregions (Figure 2), generally following IBISCH (1996). Expectedly, some of them overlap partially or to a great extent the major geographic regions listed in (1). These are:

Amazonian Rainforests: located in Beni, La Paz, Pando, as well as at the Andean foothills in Cochabamba and Santa Cruz; 100-600 m a.s.l.; mean annual temperature 29-27 °C; mean annual precipitation 1800-2200 mm,



Figure 1.- Map of Bolivian departments.
Figura 1.- Mapa de los departamentos Bolivianos.

more at the Andean foothills; 0-3 arid months; tall evergreen rainforest.

Cerrado formations: located in Santa Cruz and Beni; 200-1000 m a.s.l.; mean annual temperatures 18-25 °C; mean annual precipitation 1000-1500 mm; 3-5 arid months; mainly dry open savanna formations but includes also low dry-forests; frequent natural fires.

“Ceja de Montaña” - Cloud Forests: located in La Paz, Cochabamba, and Santa Cruz; 2500-3500 m a.s.l.; mean annual temperature 10-14 °C; mean annual precipitation 2500-3500 mm; 0-2 arid months; low evergreen cloud forest, subparamo.

Chiquitanía Forests: located in the provincias Velasco, Núflo de Chávez, Sandoval, and Chiquitos in the Departamento de Santa Cruz; forest type unique to Bolivia; transition zone between humid Amazonian forests and dry Chaco forests; 300-1200 m a.s.l.; mean annual temperatures 18-23 °C; mean annual precipitation 1000-1500 mm; 3-5 arid months; semi-deciduous forests of medium height, large parts evergreen.

Chaco Lowland Forests: located in Chuquisaca, Santa Cruz, and Tarija; 300-600 m

a.s.l.; mean annual temperature 25-26 °C; mean annual precipitation 400-900 mm, around 1000 mm at the Andean foothills and in the northern transition zone to the Pantanal; 6-8 arid months; low dry-forests with several succulent plants.

Chaco Montane Forests: located in Chuquisaca, Santa Cruz, and Tarija; 600-1500 m a.s.l.; mean annual temperatures 18-22 °C; mean annual precipitation 1000-2000 mm; 6-7 arid months; deciduous forests of medium height; relationships to Caatinga formations.

High Andean Vegetation: located in Chuquisaca, Cochabamba, La Paz, Oruro, Potosí, and Tarija; 2500-4600 m a.s.l.; *Polylepis* forest growth up to 5200 m a.s.l. around Volcán Sajama; mean annual temperature below 10 °C; mean annual precipitation 500-700 mm; 6-8 arid months; puna, low evergreen montane forests.

Humid Transition Lowland Forests: located in Beni, Cochabamba, and Santa Cruz; forest type unique to Bolivia; 150-300 m a.s.l.; mean annual temperatures around 25 °C; mean annual precipitation 1200-1800 mm; 2-4 arid months; evergreen rainforests of medium height; azonal gallery forests along rivers.

Inter-Andean Valleys: located in Chuquisaca, Cochabamba, La Paz, Santa Cruz, and Tarija; 1300-3000 m a.s.l.; mean annual temperature 12-16 °C, maximum above 30 °C, minimum below 0 °C; mean annual precipitation 500-700 mm; 6-8 arid months; deciduous and semideciduous forests of medium height; biogeographical relationships with the dry Chaco forests.

Pre-Cambrian Shield Humid Forests: located in N Santa Cruz and parts of Beni; 200-1000 m a.s.l.; mean annual temperatures 18-25 °C; mean annual precipitation 1500-1800 mm; 2-4 arid months; evergreen forest of medium height; azonal vegetation on rock outcrops (inselbergs) and sandstone ridges.

Tucumanian-Bolivian Montane Forests: located in Chuquisaca, Santa Cruz, and Tarija; 800-2600 m a.s.l.; mean annual temperature




Figure 2.- Map of Bolivian ecoregions.

Figure 2.- Mapa de las ecoregiones de Bolivia.

13-23 °C; mean annual precipitation 1000-2000 mm; 3-5 arid months; montane forests of medium height.

Wet Savannas: located in Beni, Santa Cruz, and N La Paz; 130-250 m a.s.l.; mean annual temperature around 26 °C; mean annual precipitation 1000-2000 mm; 2-6 arid months; swamps and grass savannas with scattered forest islands.

Yungas-Montane Rainforests: located in Cochabamba, La Paz, and Santa Cruz; 600-2500 m a.s.l.; divided into upper montane rainforests (1500-2500 m a.s.l.) and lower montane rainforests (600-1500 m a.s.l.); mean annual temperature 12-24 °C, minimum in upper regions below 0 °C; mean annual precipitation 2500 to more than 6000 mm, maximum rainfall between 1500 and 1800 m a.s.l.; 0-2 arid months; characterized by steep slopes and deep valleys; evergreen montane rainforest of medium height.

The transitional zones between two or more of these ecoregions may contain species of different origin (for example, in Beni, where savannas and forest intermingle to form a patchy environment). Likewise, it must be taken into account that some species (e. g., *Bufo marinus*, *Hyla geographica*, etc.) are linked to the rainforest domain, but they inhabit open areas within it (river shores, clearings, etc.). Moreover, some Cerrado and/or Chacoan amphibians can enter open areas within the rainforest domain, either as a natural colonization or, quite often, taking advantage of human perturbations to the landscape (roads, deforestation, etc.). As a result, a Chacoan or Cerrado species may be found, for example, in Pando (e. g., *Leptodactylus elenae*). Likewise, some Chacoan or Cerrado species may be found in temperate inter-Andean valleys (e. g., *Scinax squalirostris*). Finally, some Amazonian species enter the northern Chaco (e. g., *Sphaenorhynchus lacteus*, *Leptodactylus leptodactyloides*).

(4) Habitat use: Aquatic (when the species is almost always in the water), Arboreal (those

species with digital pads adapted to climb on vegetation, although they can also move on the ground), Fossorial (when the species spends a great part of its life underground, for which it has some special adaptations), and Terrestrial [many terrestrial or arboreal species are usually near bodies of water, either ponds or streams, although they do not necessarily enter the water (e. g., centrolenids)].

5) Type of habitat: Forest or Open formation. This is an important distinction. Of course, most amphibians from savannas live in open formations, but many species from the forest domain do not occur in the forest itself, but in the open areas within it (e. g., forest gaps, forest edges, river edges, human clearings, etc.). Although trees in the Chaco can be closely spaced, they are not too tall and the Chaco is thus considered an open formation. Two facts must be taken into account: first, these two kinds of major habitats are not clear-cut categories (there are open secondary forests, areas disturbed by human activity, patchy environments, etc.); and second, several species are adapted to live in transitional zones or can occur in the two types of habitats (although often not being equally abundant in both of them).

With these five kinds of information, the putative area of distribution and habitat of a particular species in the country can be easily outlined, and it is possible to predict its occurrence in areas where it has not been found before.

Key literature.-This section points out, by chronological order, the most relevant publications on each species in Bolivia, especially those that appeared after 1990. This section is not included in accounts of species reported herein for the first time, a fact which will be indicated in the "Remarks" section. Additional references, remarks, and data on the distribution of many species are found in DE LA RIVA (1990a) but, in general, this publication will not be cited again as a key literature reference for every species, unless no other rel-

event publications exist. When the species considered has its type locality in Bolivia but is not a Bolivian endemism, the original citation is provided only if it contains additional information (on taxonomy, ecology, distribution, natural history, etc.) other than the mere description of the species. When a species has been reported in the country only once, the corresponding reference is given. In some cases this publication may be a short note on distribution which, although it contains little information in general terms, represents all the information available for that particular species in the country. Papers and reports on faunal lists and surveys of particular localities or areas, inventories, etc., usually deal with no single species in particular. Thus, a paper of this kind is not mentioned in this section, unless it contains the only known Bolivian record of the species, unless the species is treated more extensively, and/or unless substantial information other than geographical is provided. Likewise, taxonomic reviews of genera or species groups which include Bolivian specimens, but where they are not the main subject, are not cited here unless they treat Bolivian endemic species. Instead, the general bibliography should be consulted.

Remarks.-This section (which does not appear if there is no particular comment about the species) includes pertinent comments on taxonomic problems, subspecies, distribution, ongoing research on particular topics, etc., with a special reference to the changes that have taken place since DE LA RIVA's (1990a) list. Recent synonymizations are indicated, especially if the synonymized species has its type locality in Bolivia, but older, well settled taxonomic changes are not usually mentioned.

In the list we have not included doubtful species [those known only from lost types, or whose type locality is unknown, etc.; see DE LA RIVA (1990a)]. However, we consider them in a separate section because future research may require us to take them into account.

The comprehensive literature on Bolivian

amphibians presented herein includes every paper in which Bolivian amphibians are considered in any way. We provide references dealing with ecology, taxonomy, distribution, etc. of Bolivian amphibian species as long as they deal with Bolivian specimens. Papers on species of amphibians that occur in Bolivia but which are based on research carried out in other countries, are not considered (for example, papers dealing with descriptions of species that occur in Bolivia but whose type locality is not in Bolivia). As an exception, a few references cited in the main text do not contain specific information on Bolivian amphibians, but are relevant to understand some problems. Besides published papers, we also list unpublished reports and dissertations. This "gray literature" is usually difficult to obtain (and we are aware that we might have overlooked some of these papers), but we consider it is important to take these papers into account because, in some instances, they deal with faunas of poorly surveyed regions and very often are the only source of information about the amphibians occurring in such areas. Communications presented at congresses and meetings are excluded, unless they were published as books or edited conference proceedings. Likewise, references in which Bolivian amphibians are treated in a vague or general context (for example, non technical books) are excluded, especially if there is more accurate information published on the same topic elsewhere. Two or more references by a given author in the same year are ordered as a, b, c, etc., only when they are cited in this way in the main text. References in the list are in alphabetical order. When there are more than two authors, the citation in the main text is given with the name of the first author followed by "*et al.*". In this case, if there are several references of the same first author and year, the citations have been marked as a, b, c, etc. in the main text, following the order with which they appear, and irrespective of the alphabetical order followed in the list of references.

RESULTS

Updated checklist of Amphibians of Bolivia.

Anura

*Bufo*idae

Atelopus tricolor Boulenger, 1902

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; LAVILLA *et al.*, 1997; LÖTTERS & DE LA RIVA, 1998; LÖTTERS *et al.*, 1999a; KÖHLER, 2000a.

Remarks: LÖTTERS & DE LA RIVA (1998) placed *Atelopus willimani* Donoso-Barros [type locality: Runerrabaque (sic) (= Rurrenabaque, Beni)] in the synonymy of *A. tricolor*.

Bufo amboroensis Harvey & Smith, 1993*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Aquatic, Terrestrial; 5) Forest.

Key literature: HARVEY & SMITH, 1993.

Bufo arenarum Hensel, 1867

Distribution: 1) Subtropical South America; 2) **Cochabamba, Chuquisaca, Santa Cruz, Tarija**; 3) Chaco Lowland Forests, Chaco Montane Forests, Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a; KÖHLER, 2000a.

Remarks: According to LAURENT (1969) the subspecies present in Bolivia is *B. arenarum arenarum*. Figure 7 in DE LA RIVA (1990a; p. 305) actually corresponds to *B. poeppigii*, as well as the record from Caranda, Santa Cruz (p. 265).

Bufo castaneoticus Caldwell, 1991

Distribution: 1) Amazonia; 2) Beni, La Paz,

Pando; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Key literature: KÖHLER & LÖTTERS, 1999a.

Bufo fissipes Boulenger, 1903

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1990a.

Remarks: FROST (1985) reported this Peruvian species (type locality: Santo Domingo, Puno) in the Departamento of Cochabamba, but we are unaware of voucher specimens other than the holotype. It is surprising that several species of the *B. veraguensis* group have been described in the last years from Peru and Bolivia (and some others still await description), but this species has yet to be collected in Bolivia..

Bufo granulosus Spix, 1824

Distribution: 1) Middle and South America; 2) **Beni, Cochabamba, Chuquisaca, La Paz, Pando, Santa Cruz, Tarija**; 3) Amazonian Rainforests, Cerrado, Chiquitania Forests, Chaco Lowland Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: REYNOLDS & FOSTER, 1992; KÖHLER, 1995a; REICHLE, 1996; 1997a; KÖHLER *et al.*, 1997; HARVEY *et al.*, 1998; REICHLE & KÖHLER, 1998.

Remarks: *Bufo granulosus* is a polytypic species with 14 described subspecies (GALLARDO, 1965). Some of them have been considered as full species by different authors. We agree that some of these taxa deserve specific status, but a revision of the whole complex is desirable before any taxonomic decision is made. Four subspecies have been reported for Bolivia: *B. g. goeldii* Gallardo in the north of the Bolivian Amazon, *B. g. major* Müller & Hellmich in the Chacoan area (type locality: San José de Chiquitos, Santa Cruz), *B. g. mini* Gallardo in

the Beni area, (type locality: Upper Beni, below mouth of Mapiri) and *B. g. mirandaribeiroi* Gallardo in northern Santa Cruz. *Bufo g. goeldii*, *B. g. mini*, and *B. g. major* are similar in morphology, although *B. g. major* is larger than the two other subspecies. The most distinctive form is *B. g. mirandaribeiroi*, which otherwise seems to be more closely related to the nominal subspecies (from NE Brazil) than the other Bolivian subspecies. Thus, at least two different lineages seem to be present in the country. In summary, the splitting of *B. granulosus* might lead to the recognition of 2-4 species in Bolivia.

***Bufo guttatus* Schneider, 1799**

Distribution: 1) Amazonia; 2) Beni, **La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: LÖTTERS *et al.*, 2000.

Bufo justinianoi* Harvey & Smith, 1994*

Distribution: 1) CFP; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: HARVEY & SMITH, 1994; KÖHLER, 2000a.

***Bufo "margaritifer"* (Laurenti, 1768)**

Distribution: 1) Tropical Middle and South America; 2) **Beni, Cochabamba, Chuquisaca, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Inter-Andean Valleys, Pre-Cambrian Shield Humid Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest, Open.

Key literature: REYNOLDS & FOSTER, 1992; KÖHLER, 1995a; DE LA RIVA *et al.*, 1996a; KÖHLER *et al.*, 1997; KÖHLER & LÖTTERS, 1999a.

Remarks: The systematics of the *B. margari-*

tifer (or *B. typhonius*) complex is perhaps one of the most chaotic of all Neotropical anuran groups (HOOGMOED, 1986; 1989; 1990; HASS *et al.*, 1995). HOOGMOED (1986) stated that the small toads of the Bolivian lowlands are similar to *B. acutirostris* Spix which otherwise occurs in E Colombia, S Venezuela and adjacent Brazil. Bioacoustic studies carried out in N Santa Cruz, northern Chaco of Santa Cruz, and Beni indicate that all these populations are conspecific (DE LA RIVA *et al.*, 1996a; KÖHLER *et al.*, 1997). KÖHLER & LÖTTERS (1999a) stated that *B. castaneoticus* is one of two forms that occur in the Departament of Pando, and LÖTTERS & KÖHLER (2000a) are describing a new species from the humid montane forests of Cochabamba and Santa Cruz. The type locality of *Bufo pleuropterus* Schmidt (currently considered a synonym of *B. margaritifer*) is given as “Grenzgebiet von Bolivia gegen Peru, in etwa 3000 [feet] Höhe” (at the border of Bolivia and Peru, approximately 700 m) by SCHMIDT (1858). According to the loss of Bolivian territory in 1909 and the travel route of the collector J. v. Warszewiez, this locality is most probably situated in present-day Peru. The taxonomic status of some populations of the *B. margaritifer* complex in the Bolivian lowlands and valleys remains a topic to be investigated, but it is almost sure that not less than four different species occur in the country.

***Bufo marinus* (Linnaeus, 1758)**

Distribution: 1) Middle and South America; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA *et al.*, 1996a.

***Bufo paracnemis* Lutz, 1925**

Distribution: 1) Bolivian-Argentinean Forests, Cerrado, Chaco, Subtropical South America; 2) **Beni, Cochabamba, Chuquisaca, Santa**

Cruz, Tarija; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas; 4) Terrestrial; 5) Open. Key literature: REYNOLDS & FOSTER, 1992; DE LA RIVA, 1993a; KÖHLER, 1995a; 2000a; REICHLE, 1996; 1997a; KÖHLER *et al.*, 1997.

Bufo poeppigii Tschudi, 1845

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz;** 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Terrestrial. 5) Forest, Open.

Key literature: DE LA RIVA *et al.*, 1996a; KÖHLER, 2000a.

Remarks: See remarks under *B. arenarium*.

Bufo quechua Gallardo, 1961**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz;** 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: GALLARDO, 1961a; HARVEY & SMITH, 1993; 1994; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995a.

Remarks: HARVEY & SMITH (1993) placed *Bufo echinodes* Reynolds & Foster [type locality: Parjacti (sic), Cochabamba (=Paracuti)] in the synonymy of *B. quechua*.

Bufo spinulosus Wiegmann, 1834

Distribution: 1) Andean Highlands; 2) **Cochabamba, Chuquisaca, La Paz, Oruro, Potosí;** 3) High Andean Vegetation, Inter-Andean Valleys; 4) Terrestrial.

Key literature: ERGUETA, 1991; APARICIO, 1993; IBISCH & BÖHME, 1993; KÖHLER, 1995a.

Remarks: GALLARDO (1961a) described the subspecies *B. s. altiperuvianus* from Bolivia (type locality: Challapata, Oruro). HARVEY (1997) reported a toad tentatively identified as *B. gallardoi* Carrizo from El Palmar (Chuquisaca). This poorly known Argen-

tinean species might be related either to the *spinulosus* group (CARRIZO, 1992), or to the *marinus* group (HARVEY, 1997). Although *B. gallardoi* or a similar form most probably occurs in the country, we do not include it in the list until a reliable record exists.

Bufo veraguensis Schmidt, 1857

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, Chuquisaca, La Paz, Santa Cruz, Tarija;** 3) Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; HARVEY & SMITH, 1993; 1994; KÖHLER *et al.*, 1995b; KÖHLER, 2000a.

Remarks: Some species in the *B. veraguensis* group remain to be described, whereas the taxonomic status of others is doubtful. See remarks under *B. fissipes*.

Dendrophryniscus minutus (Melin, 1941)

Distribution: 1) Amazonia; 2) Beni, **Cochabamba, La Paz, Pando, Santa Cruz;** 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1999a.

Melanophryniscus rubriventris (Vellard, 1947)

Distribution: 1) Temperate Inter-Andean Valleys; 2) **Cochabamba, Chuquisaca, Santa Cruz, Tarija;** 3) Inter-Andean Valleys; 4) Terrestrial.

Key literature: DE LA RIVA, 1995a.

Remarks: The Bolivian specimens known seem to belong to the nominotypical form.

Centrolenidae

Cochranella bejaranoi (Cannatella, 1980)**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, Chuquisaca, La Paz, Santa Cruz;** 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: CANNATELLA, 1980; CANNATELLA & DUELLMAN, 1982; SANCHÍZ & DE

LA RIVA, 1993; HARVEY, 1996; MÁRQUEZ *et al.*, 1996; KÖHLER, 2000a.

Remarks: HARVEY (1996) placed *Centrolenella flavidigitata* Reynolds & Foster (type locality: Road to San Onofre, 3.3 km by road N of the road from Cochabamba to Villa Tunari, Cochabamba) in the synonymy of *C. bejaranoi*.

Cochranella nola Harvey, 1996**

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) Cochabamba, Chuquisaca, **Santa Cruz**; 3) Amazonian Rainforests, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: HARVEY, 1996; KÖHLER, 2000a; LÖTTERS & KÖHLER, 2000b.

Cochranella pluvialis (Cannatella & Duellman, 1982)

Distribution: 1) Humid Montane Forests; 2) **La Paz**, Cochabamba; 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: KÖHLER & REICHLE, 1998.

Hyalinobatrachium bergeri (Cannatella, 1980)*

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: CANNATELLA, 1980; CANNATELLA & DUELLMAN, 1982; SANCHÍZ & DE LA RIVA, 1993; MÁRQUEZ *et al.*, 1996; KÖHLER, 2000a.

Dendrobatidae

Allobates femoralis (Boulenger, 1884 “1883”)

Distribution: 1) Amazonia; 2) Beni, **La Paz, Pando**; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Remarks: Found by J. E. Cadle and S. Reichle in November 1999 and reported herein for the first time.

Colostethus brunneus (Cope, 1887)

Distribution: 1) Amazonia, Cerrado; 2) **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: GONZÁLES *et al.*, 1999.

Colostethus mcdiarmidi Reynolds & Foster, 1992**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; GONZÁLES *et al.*, 1999.

Colostethus trilineatus (Boulenger, 1884 “1883”)

Distribution: 1) Southwestern Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Terrestrial; 5) Forest.

Key literature: MORALES, 1994; DE LA RIVA *et al.*, 1996b; REICHLE & KÖHLER, 1996a; GONZÁLES *et al.*, 1999; KÖHLER & LÖTTERS, 1999a.

Remarks: MORALES (1994) stated that *C. trilineatus* is the proper name for southern populations of Amazonian *Colostethus* formerly considered as *C. marchesianus* (Melin). Populations not formally described from Cochabamba (EDWARDS, 1974) are also *C. trilineatus* (DE LA RIVA *et al.*, 1996a). An unidentified, related species occurs in Pando (KÖHLER & LÖTTERS, 1999a).

Epipedobates boliviensis (Boulenger, 1902)**

Distribution: 1) Humid Montane Forests; 2) **La Paz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: BOULENGER, 1902; SILVERSTONE, 1976; RODRÍGUEZ & MYERS, 1993; GONZÁLES *et al.*, 1999.

Remarks: This species has been only recently rediscovered after almost one century

without new findings (GONZÁLES *et al.*, 1999).

***Epipedobates hahneli* (Boulenger, 1884 “1883”)**

Distribution: 1) Upper Amazon Basin; 2) Beni, **La Paz, Pando**; 3) Amazonian Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: DE LA RIVA *et al.*, 1996b; GONZÁLES *et al.*, 1999; KÖHLER & LÖTTERS, 1999a.

Epipedobates pictus* (Bibron in Tschudi, 1838)

Distribution: 1) Southwestern Amazonia, Cerrado, Humid Montane Forests; 2) **Beni, Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest, Open.

Key literature: REYNOLDS & FOSTER, 1992; DE LA RIVA, 1993a; HADDAD & MARTINS, 1994; KÖHLER, 1995a; 2000a; DE LA RIVA *et al.*, 1996b; KÖHLER & LÖTTERS, 1999a; GONZÁLES *et al.*, 1999.

Remarks: An undescribed species from La Paz similar to *E. pictus* is being studied by S. Lötters and S. Reichle.

***Epipedobates trivittatus* (Spix, 1824)**

Distribution: 1) Amazonia; 2) Beni, La Paz, **Pando**; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Remarks: Found by J. E. Cadle and S. Reichle in November 1999 and reported herein for the first time.

Hylidae

Hemiphractinae

Gastrotheca lauzuricae* De la Riva, 1992 “1991” *

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

ñña-Cloud Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1992a.

Remarks: Known only from the holotype.

***Gastrotheca marsupiata* (Duméril & Bibron, 1841)**

Distribution: 1) Andean Highlands, Humid Montane Forests; 2) **Cochabamba, Chuquisaca, La Paz, Oruro, Potosí, Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, High Andean Vegetation, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest, Open.

Key literature: ERGUETA, 1991; DE LA RIVA, 1992a; VELLARD, 1992; APARICIO, 1993; IBISCH & BÖHME, 1993; DE LA RIVA *et al.*, 1995; KÖHLER, 1995a; KÖHLER *et al.*, 1995a.

Remarks: The variation in populations from the highlands and those from the Yungas was studied by DE LA RIVA (1992a), who suggested the possibility that more than one species is hidden under the name *G. marsupiata* in Bolivia.

***Gastrotheca splendens* (Schmidt, 1857)**

Distribution: 1) Humid Montane Forests; 2) Cochabamba, **Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1992a; DUELLMAN & DE LA RIVA, 1999.

Remarks: This species was “lost” for almost a century and a half. The type locality is unknown, and the two only specimens known at present are the holotype and a specimen recently collected in cloud forests of Santa Cruz (DUELLMAN & DE LA RIVA, 1999). It is likely that *G. splendens* is a Bolivian endemism.

***Gastrotheca testudinea* (Jiménez de la Espada, 1871)**

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1992a; KÖHLER, 2000a.

Hylineae

Hyla acreana Bokermann, 1964

Distribution: 1) Southwestern Amazonia; 2)

Beni, Cochabamba, La Paz, Pando, Santa Cruz; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993.

Hyla albonigra Nieden, 1923**

Distribution: 1) Andean Highlands, Temperate Inter-Andean Valleys; 2) **Cochabamba, Chuquisaca, Potosí, Santa Cruz;** 3) High Andean Vegetation, Inter-Andean Valleys; 4) Arboreal; 5) Open.

Key literature: DUMÉRIL & BIBRON, 1841; D'ORBIGNY, 1847; NIEDEN, 1923; ANDERSSON, 1938; DUELLMAN *et al.*, 1997.

Remarks: DUELLMAN *et al.* (1997) placed *Hyla ocapia* Andersson (type locality: Puka Khara, Potosí) in the synonymy of *H. albonigra*.

Hyla albopunctata Spix, 1824

Distribution: 1) Cerrado; 2) **Santa Cruz;** 3) Cerrado; 4) Arboreal; 5) Open.

Key literature: DE SÁ, 1995; DE LA RIVA *et al.*, 1997.

Hyla andina Müller, 1924

Distribution: 1) Andean Highlands, Humid Montane Forests, Temperate Inter-Andean Valleys; 2) **Cochabamba, Chuquisaca, La Paz, Santa Cruz, Tarija;** 3) Ceja de Montaña-Cloud Forests, High Andean Vegetation, Inter-Andean Valleys, Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest, Open.

Key literature: ERGUETA, 1991; REYNOLDS & FOSTER, 1992; MÁRQUEZ *et al.*, 1993; KÖH-

LER, 1995a; 2000a; KÖHLER *et al.*, 1995b; DUELLMAN *et al.*, 1997.

Remarks: DUELLMAN *et al.* (1997) elevated *Hyla pulchella andina* to specific status. *Hyla andina* shows remarkable variation and further studies might demonstrate that more than one species is involved. Figures 5 and 6 in KÖHLER *et al.* (1995b; p. 17), purportedly showing specimens of *H. pulchella andina*, actually correspond to *H. marianitae*. HARVEY (1997) was unsure if specimens he assigned to *H. pulchella* (= *H. andina*) from El Palmar (Chuquisaca) could actually be *Hyla varelae* Carrizo. He remarked that Carrizo's description was inadequate to make a definite identification without examination of comparative material.

Hyla armata Boulenger, 1902*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz;** 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; MÁRQUEZ *et al.*, 1993; DUELLMAN *et al.*, 1997; KÖHLER, 2000a.

Remarks: There is considerable variation in skin texture, color pattern, color of the iris and, apparently, advertisement calls, which suggests that more than one species might be involved under the name *H. armata* (DUELLMAN *et al.*, 1997).

Hyla balzani Boulenger, 1898*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz;** 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; DUELLMAN *et al.*, 1997; KÖHLER, 2000a.

Remarks: DUELLMAN *et al.* (1997) placed *Hyla callipleura* Boulenger [type locality: Charuplaya (Cochabamba) and San Ernesto (La Paz)] in the synonymy of *H. balzani*. However, further evidence based mainly on vocalizations seems to indicate that perhaps *H. callipleura* should be resurrected (J. Köhler

and S. Reichle, pers. obs.). Additionally, some frog populations of the *H. pulchella* group in the inter-Andean temperate valleys possibly represent an undescribed species (see KÖHLER *et al.*, 1995b; DUELLMAN *et al.*, 1997).

Hyla bifurca Andersson, 1945

Distribution: 1) Upper Amazon Basin; 2) Beni, Cochabamba, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993.

Remarks: There are considerable differences in color pattern between Santa Cruz populations and those from Ecuador (where the species has its type locality). In Pando there are specimens like those from Ecuador and those from Santa Cruz, as well as intermediate ones (DE LA RIVA, 1993a).

Hyla boans (Linnaeus, 1758)

Distribution: 1) Tropical Middle and South America; 2) Beni, Cochabamba, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1990a.

Hyla calcarata Troschel, 1848

Distribution: 1) Upper Amazon Basin; 2) **La Paz**, **Pando**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1990a; APARICIO, 1992.

Hyla charazani Vellard, 1970**

Distribution: 1) Temperate Inter-Andean Valleys; 2) **La Paz**; 3) Ceja de Montaña-Cloud Forests, Inter-Andean Valleys; 4) Aquatic, Arboreal; 5) Open.

Key literature: VELLARD, 1970; DUELLMAN *et al.*, 1997.

Remarks: This poorly known species (see DUELLMAN *et al.*, 1997, p. 12) was recently rediscovered by J. Aparicio and S. Reichle at the type locality. It is similar to *Hyla armata*, but seems to represent a different species whose status must be studied together with that of certain populations currently considered as *H. armata* (see remarks on *H. armata*).

Hyla chlorostea Reynolds & Foster, 1992**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba**; 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992.

Remarks: Known only from the holotype. Uncertain taxonomic status.

Hyla fasciata Günther, 1859

Distribution: 1) Upper Amazon Basin; 2) **Beni**, **Cochabamba**, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993.

Remarks: DE LA RIVA (1990a) placed *Hyla steinbachi* Boulenger (type locality: Province Sarah, Santa Cruz) in the synonymy of *H. fasciata*.

Hyla geographica Spix, 1824

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: DE LA RIVA's (1990a) statement that the subspecies present in Bolivia is *H. g. punctatissima* (Reinhardt & Lütken) is unfounded.

Hyla granosa Boulenger, 1882

Distribution: 1) Amazonia; 2) La Paz, **Pando**; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest, Open.

Key literature: KÖHLER & LÖTTERS, 1999a.

Hyla koechlini Duellman & Trueb, 1989

Distribution: 1) Southwestern Amazonia; 2) **Beni**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Remarks: This is the first record for the species in the country.

Hyla lanciformis (Cope, 1870)

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: MÁRQUEZ *et al.*, 1993.

Hyla leali Bokermann, 1964

Distribution: 1) Southwestern Amazonia; 2) **Beni**, **Cochabamba**, La Paz, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: This species was erroneously considered a junior synonym of *Hyla rossalleni* Goin by DUELLMAN (1974). Although HEYER (1977) clearly stated that both species were different and further publications treated *H. leali* and *H. rossalleni* as separate species [for example, FROST (1985)], a certain degree of confusion persisted for many years. This situation led DE LA RIVA (1990a) to report *H. rossalleni* in Bolivia. Finally, DE LA RIVA & DUELLMAN (1997) clarified the taxonomic status of this last mentioned species, which is easily distinguishable from *H. leali* and belongs to a different species

group (the *H. leucophyllata* group). Although *H. rossalleni* has not been found in Bolivia, its occurrence in the country is expected.

Hyla leucophyllata (Beireis, 1783)

Distribution: 1) Amazonia; 2) **Beni**, Cochabamba, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Hyla marianniae Carrizo, 1992

Distribution: 1) Bolivian-Argentinean Forests, Humid Montane Forests, Temperate Inter-Andean Valleys; 2) **Cochabamba**, **Chuquisaca**, **Santa Cruz**, Tarija; 3) Amazonian Rainforests, Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest, Open.

Key literature: MÁRQUEZ *et al.*, 1993; DUELLMAN *et al.*, 1997; HARVEY, 1997; LÖTTERS *et al.*, 1999b; KÖHLER, 2000a.

Remarks: The description of the call of *H. callipleura* (= *H. balzani*) in MÁRQUEZ *et al.* (1993) actually corresponds to *H. marianniae*, as stated by DUELLMAN *et al.* (1997). See remarks under *H. andina*.

Hyla marmorata (Laurenti, 1768)

Distribution: 1) Upper Amazon Basin; 2) **Beni**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: KÖHLER, 1995a; 1995b; KÖHLER & BÖHME, 1996.

Hyla melanargyrea Cope, 1887

Distribution: 1) Cerrado; 2) **Santa Cruz**; 3) Cerrado, Chiquitanía Forests; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; KÖHLER & BÖHME, 1996.

Hyla minuta Peters, 1872

Distribution: 1) South America; 2) **Beni**, Cochabamba, **Chuquisaca**, La Paz, **Pando**, **Santa Cruz**, **Tarija**; 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Inter-Andean Valleys, Pre-Cambrian Shield Humid Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: A species related to *H. minuta* from the Chapare region of Cochabamba is being described by KÖHLER & LÖTTERS (2000).

Hyla nana Boulenger, 1889

Distribution: 1) Cerrado, Chaco, Subtropical South America; 2) **Beni**, Chuquisaca, **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: LANGONE & BASSO, 1987; DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Hyla parviceps Boulenger, 1882

Distribution: 1) Upper Amazon Basin; 2) Beni, Cochabamba, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993.

Hyla punctata (Schneider, 1799)

Distribution: 1) Tropical South America; 2) **Beni**, Cochabamba, **La Paz**, **Pando**, **Santa**

Cruz; 3) Amazonian Rainforests, Chiquitanía Forests, Chaco Lowland Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: The validity of the Chacoan subspecies *Hyla punctata rubrolineata* Lutz (type locality: Buenavista, Santa Cruz) is doubtful.

Hyla raniceps (Cope, 1862)

Distribution: 1) Cerrado, Chaco; 2) **Beni**, **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Hyla rhodopepla Günther, 1859

Distribution: 1) Upper Amazon Basin; 2) Beni, **Cochabamba**, **La Paz**, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest, Open.

Key literature: MÁRQUEZ *et al.*, 1993.

Hyla riveroi Cochran & Goin, 1970

Distribution: 1) Upper Amazon Basin; 2) **Beni**, **Cochabamba**, **La Paz**, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1993; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Hyla rubricundula Reinhardt & Lütken, 1862

Distribution: 1) Cerrado; 2) **Santa Cruz**; 3) Cerrado, Chiquitanía Forests; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1990a.

Hyla sarayacuensis Shreve, 1935

Distribution: 1) Upper Amazon Basin; 2) Beni, La Paz, **Pando**; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Remarks: This is the first record for the species in the country.

Hyla schuberti Bokermann, 1963

Distribution: 1) Southwestern Amazonia; 2) Beni, **La Paz**, **Pando**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1999b.

Hyla tritaeniata Bokermann, 1965

Distribution: 1) Cerrado; 2) **Beni**, **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1996a; 1998.

Remarks: Based on this species' remarkable intra- and interpopulational variation, NAPOLI & CARAMASCHI (1999) suggested that *H. tritaeniata* is a complex of species.

Osteocephalus buckleyi (Boulenger, 1882)

Distribution: 1) Upper Amazon Basin; 2) **Beni**, **Cochabamba**, La Paz, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1990a.

Osteocephalus leprieurii (Duméril & Bibron, 1841)

Distribution: 1) Upper Amazon Basin; 2) Beni, La Paz, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1995.

Remarks: SMITH *et al.* (2000) are describing a new species of Bolivian *Osteocephalus* which corresponds to what is here considered *O. leprieurii*.

Osteocephalus pearsoni (Gaige, 1929)*

Distribution: 1) Southwestern Amazonia; 2) **Beni**, Cochabamba, **La Paz**, Pando; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Key literature: GAIGE, 1929; DE LA RIVA, 1990a.

Osteocephalus taurinus Steindachner, 1862

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1995; KÖHLER, 1995a; KÖHLER & BÖHME, 1996.

Phrynohyas coriacea (Peters, 1867)

Distribution: 1) Upper Amazon Basin; 2) Beni, La Paz, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; 1994a; DE LA RIVA *et al.*, 1995.

Phrynohyas resinifictrix (Goeldi, 1907)

Distribution: 1) Amazonia; 2) Beni, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Remarks: Found by J. E. Cadle and S. Reichle in November 1999 and reported here for the first time.

Phrynohyas venulosa (Laurenti, 1768)

Distribution: 1) Middle and South America; 2) **Beni**, **Cochabamba**, **Chuquisaca**, **La Paz**, **Pando**, **Santa Cruz**, Tarija; 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Inter-Andean Valleys, Pre-Cambrian Shield Humid Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; DE LA

RIVA *et al.*, 1995; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Scarthyla ostinodactyla Duellman & de Sá, 1988

Distribution: 1) Upper Amazon Basin; 2) La Paz, **Pando**; 3) Amazonian Rainforests; 4) Aquatic, Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1999c.

Scinax acuminatus (Cope, 1862)

Distribution: 1) Cerrado, Chaco; 2) Chuquisaca, **Santa Cruz**, Tarija; 3) Cerrado, Chaco Lowland Forests, Wet Savannas; 4) ArboREAL; 5) Open.

Key literature: DE LA RIVA, 1990a.

Scinax castroviejoi De la Riva, 1993*

Distribution: 1) Bolivian-Argentinean Forests, Temperate Inter-Andean Valleys; 2) Cochabamba, Chuquisaca, La Paz, **Santa Cruz**, Tarija; 3) Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) ArboREAL; 5) Open.

Key literature: DE LA RIVA, 1993b; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b.

Remarks: This species was originally described from temperate valleys of Santa Cruz and from a locality in Jujuy (Argentina). Subsequent surveys carried out at the Jujuy locality failed to find the species. Instead, *S. fuscovarius* was common. These are sibling species difficult to distinguish morphologically, but they have very different advertisement calls. Argentinean paratypes of *S. castroviejoi* were assigned to this species based on morphology only. Thus, it is possible that these specimens actually are *S. fuscovarius*, a species whose geographic variation could have enhanced the similarity between the Jujuy and Santa Cruz populations. Bolivian samples of these two species can be distinguished by meristic characters, but perhaps this is not the case

when other populations are considered. However, many species have a distribution comprising the temperate and subtropical valleys of NW Argentina up to Santa Cruz or Cochabamba (HARVEY, 1997). Thus, since the presence of one species does not exclude the other (they are sympatric at some Bolivian localities), the occurrence of *S. castroviejoi* in Argentina would still be plausible even if the paratypes from Jujuy proved to be *S. fuscovarius*.

Scinax chiquitanus (De la Riva, 1990)*

Distribution: 1) Southwestern Amazonia; 2) **Beni**, Cochabamba, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) ArboREAL; 5) Forest.

Key literature: DE LA RIVA, 1990b; 1993a; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a.

Remarks: DUELLMAN & WIENS (1993) provided data on this species from Peru. The male shown in their Fig. 2 (p. 8) has a color pattern which differs markedly from that of Bolivian males, which otherwise present little variation. The taxonomic status of Peruvian specimens identified as *S. chiquitanus* should be reassessed.

Scinax fuscovarius (Lutz, 1925)

Distribution: 1) Cerrado, Chaco; 2) **Beni**, Chuquisaca, **Santa Cruz**, Tarija; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Inter-Andean Valleys, Pre-Cambrian Shield Humid Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas; 4) Terrestrial, ArboREAL; 5) Open.

Key literature: DE LA RIVA, 1993a; 1993b; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a; 2000a; KÖHLER & BÖHME, 1996.

Remarks: We follow LUTZ (1973) and CEI (1980) in considering *Scinax trachythorax* (Müller & Hellmich) as a synonym of *S. fuscovarius*.

Scinax garbei (Miranda-Ribeiro, 1926)

Distribution: 1) Upper Amazon Basin; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Cerrado, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1994.

Remarks: DUELLMAN & WIENS (1993) placed *S. epacrorthina* (Duellman) [which was reported for Bolivia; see DE LA RIVA (1990a)] in the synonymy of *S. garbei*.

Scinax nasicus (Cope, 1862)

Distribution: 1) Cerrado, Chaco; 2) Beni, Cochabamba, Chuquisaca, **Santa Cruz**, Tarija; 3) Cerrado, Chiquitanía Forests; 4) Arboreal, Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993b; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a.

Scinax nebulosus (Spix, 1824)

Distribution: 1) Lower Amazon Basin; 2) Beni, **Santa Cruz**; 3) Amazonian Rainforests, Cerrado, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1990a; HOOGMOED, 1993; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a.

Scinax parkeri (Gaige, 1929)*

Distribution: 1) Lower Amazon Basin, Cerrado; 2) **Beni, Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA *et al.*, 1994; 1997; KÖHLER, 1995a; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: Many authors have followed LUTZ (1973) in considering this species a junior synonym of *S. fuscomarginatus* (Lutz), but comparisons of the calls of both species car-

ried out by DE LA RIVA *et al.* (1994; 1997) suggested that they are not conspecific. However, this must be corroborated with more data. As stated by DE LA RIVA *et al.* (1997), the taxonomic status of the Amazonian species of the *Scinax staufferi* group needs urgent revision. In Bolivia, at least another species (probably undescribed) in the *staufferi* group occurs in the open areas of Beni and Santa Cruz.

Scinax ruber (Laurenti, 1768)

Distribution: 1) Tropical Middle and South America; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1994; KÖHLER, 1995a; KÖHLER & BÖHME, 1996; REICHLE, 1996; 1997b; REICHLE & KÖHLER, 1998.

Remarks: Given the broad distribution and variation of *S. ruber*, probably there are more than one species hidden under this name (DUELLMAN & WIENS, 1993). As far as we know, nobody is currently reviewing this species complex. Given that the type locality (Paramaribo, Surinam) is far away from Bolivia, it would be likely that Bolivian populations would receive another name if *S. ruber* were split into several species.

Scinax squalirostris (A. Lutz, 1925)

Distribution: 1) Cerrado; 2) Beni, **La Paz, Santa Cruz**; 3) Cerrado; 4) Arboreal; 5) Open.

Remarks: This species is reported here for the first time, based on specimens collected by J. Bosch, I. De la Riva and S. Reichle near Apolo, La Paz, in November 1999. The occurrence of *S. squalirostris* in the country was expected in the Pantanal area of Santa Cruz. Admittedly, its discovery for the first time at a locality so far away from the Pan-

tanal is surprising. However, the temperate valleys around Apolo are suitable for this and other species typical of temperate lowlands and/or valleys (e. g., *Scinax castroviejoi* and *Phyllomedusa boliviana*, which were also found in the area). We predict that other populations of *S. squalirostris* will be found in suitable habitats of La Paz, Beni and Santa Cruz.

Sphaenorhynchus lacteus (Daudin, 1802)

Distribution: 1) Amazonia; 2) **Beni**, Cochabamba, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1995; KÖHLER, 1995a; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Phyllomedusinae

Phyllomedusa bicolor (Boddaert, 1772)

Distribution: 1) Upper Amazon Basin; 2) **Beni**, La Paz, **Pando**, Santa Cruz; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Key literature: REICHLE *et al.*, 1997.

Phyllomedusa boliviana Boulenger, 1902*

Distribution: 1) Bolivian-Argentinean Forests, Cerrado, Humid Montane Forests, Temperate Inter-Andean Valleys; 2) **Beni**, Cochabamba, **Chuquisaca**, **La Paz**, **Santa Cruz**, **Tarija**; 3) Chiquitanía Forests, Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Open.

Key literature: CANNATELLA, 1983; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; DE LA RIVA, 2000.

Remarks: CANNATELLA (1983) synonymized *Phyllomedusa pailona* Shreve (type locality: El Pailón, Santa Cruz) with *P. boliviana* and stated that FUNKHouser's (1957) pur-

ported specimens of *P. boliviana* actually corresponded to a different species (see remarks under *P. camba*).

Phyllomedusa camba De la Riva, 2000 "1999"*

Distribution: 1) Southwestern Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1995; KÖHLER & LÖTTERS, 1999a; DE LA RIVA, 2000.

Remarks: The existence of this species, confused with *P. boliviana* by several authors, was announced long ago by CANNATELLA (1983) and it has been repeatedly reported (usually as *Phyllomedusa* sp.) from several localities in SW Peru, Brazil, and Bolivia (see DE LA RIVA, 2000).

Phyllomedusa hypochondrialis (Daudin, 1800)

Distribution: 1) Tropical South America; 2) **Beni**, **Santa Cruz**; 3) Chiquitanía Forests, Chaco Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA *et al.*, 1995; KÖHLER, 1995a; REICHLE, 1996; 1997c.

Remarks: LESCURE *et al.* (1995) stated that the correct spelling of the specific name is *hypochondrialis* instead *hypocondrialis*, a name frequently found in the literature. The record of *P. rohdei* Mertens in Beni by FUNKHouser (1957) must be in error and correspond to *P. hypochondrialis*. After CEI (1980), the subspecies *P. h. azurea* Cope occurs in the Chaco, whereas the nominal subspecies would occur in N South America. The distribution of both subspecies seems to be disjunct. The variation and taxonomic status of Bolivian populations remain to be studied.

Phyllomedusa palliata Peters, 1872

Distribution: 1) Upper Amazon Basin; 2) **Beni**, Cochabamba, La Paz, **Pando**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Arboreal; 5) Forest.
Key literature: REICHLE & KÖHLER, 1996a; KÖHLER & LÖTTERS, 1999a.

Phyllomedusa sauvagii Boulenger, 1882

Distribution: 1) Chaco; 2) Chuquisaca, **Santa Cruz**, **Tarija**; 3) Chaco Lowland Forests; 4) Arboreal; 5) Open.

Key literature: DE LA RIVA, 1990a; GONZÁLES, 1998.

Phyllomedusa tomopterna (Cope, 1868)

Distribution: 1) Upper Amazon Basin; 2) Beni, Cochabamba, **La Paz**, Pando; 3) Amazonian Rainforests; 4) Arboreal; 5) Forest.

Remarks: This is the first record for the species in the country.

Phyllomedusa vaillanti Boulenger, 1882

Distribution: 1) Upper Amazon Basin; 2) Beni, **Cochabamba**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1995.

Leptodactylidae

Ceratophryinae

Ceratophrys cornuta (Linnaeus, 1758)

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, La Paz, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Fossorial, Terrestrial; 5) Forest.
Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1995.

Ceratophrys cranwelli Barrio, 1980

Distribution: 1) Chaco; 2) Chuquisaca, **Santa**

Cruz, **Tarija**; 3) Chiquitanía Forests, Chaco Lowland Forests; 4) Fossorial, Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a; GONZÁLES, 1998.

Chacophrys pierottii (Vellard, 1948)

Distribution: 1) Chaco; 2) Chuquisaca, Santa Cruz, **Tarija**; 3) Chaco Lowland Forests; 4) Fossorial, Terrestrial; 5) Open.

Key literature: KÖHLER, 1997a.

Lepidobatrachus laevis Budgett, 1899

Distribution: 1) Chaco; 2) Chuquisaca, **Santa Cruz**, **Tarija**; 3) Chaco Lowland Forests; 4) Aquatic, Fossorial, Terrestrial; 5) Open.

Key literature: KÖHLER, 1997b.

Leptodactylinae

Adenomera andreae Müller, 1923

Distribution: 1) Amazonia; 2) Beni, **Cochabamba**, La Paz, **Pando**, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1995b; MÁRQUEZ *et al.*, 1995.

Adenomera diptyx (Boettger, 1885)

Distribution: 1) Cerrado; 2) Chuquisaca, **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1995b; MÁRQUEZ *et al.*, 1995; DE LA RIVA, 1996.

Remarks: This species is very similar to *A. hylaedactyla* but differs in its reproductive mode (DE LA RIVA, 1995b). It enters the humid Chaco. Its advertisement call was described by MÁRQUEZ *et al.* (1995) as that of *A. andreae*.

Adenomera hylaedactyla (Cope, 1868)

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, **Pando**, **Santa Cruz**; 3) Ama-

zonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1995b; KÖHLER, 1995a; 2000a; MÁRQUEZ *et al.*, 1995; KÖHLER & BÖHME, 1996; REICHLE, 1997c.

Remarks: See remarks under *A. diptyx*.

Leptodactylus boliviensis Boulenger, 1898*

Distribution: 1) Tropical Middle and South America; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1995.

Remarks: *Leptodactylus insularum* Barbour from northern South America and Middle America has been often considered a synonym of *L. boliviensis*. If it proved to be a different species, then the corresponding pattern of distribution of *L. boliviensis* would be Amazonian, instead of Tropical Middle and South America.

Leptodactylus bufonius Boulenger, 1894

Distribution: 1) Cerrado, Chaco; 2) **Chuquisaca, Santa Cruz, Tarija**; 3) Chaco Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: HEYER, 1978; DE LA RIVA, 1990a.

Leptodactylus chaquensis Cei, 1950

Distribution: 1) Cerrado, Chaco; 2) **Beni, Cochabamba, Chuquisaca, La Paz, Pando, Santa Cruz, Tarija**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a; KÖHLER, 1995a; 2000a; REICHLE, 1996; 1997c; KÖHLER & LÖTTERS, 1999a; DE LA RIVA & MALDONADO, 1999.

Remarks: As stated by DE LA RIVA & MALDONADO (1999), with the present state of knowledge *L. chaquensis* is indistinguishable from *L. macrosternum* Miranda-Ribeiro, which is distributed from N South America to Brazil and Bolivia (FROST, 1985). The only revision of the *L. ocellatus* group was carried out by GALLARDO (1964a), who reported *L. chaquensis* and *L. macrosternum* in Bolivia, but it is impossible to assign these species to localities with certainty. Provisionally, it can be assumed that Amazonian records (Beni, Cochabamba, La Paz, northern Santa Cruz) correspond to *L. macrosternum*, whereas *L. chaquensis* is rather restricted to the Chaco (Chuquisaca, S Santa Cruz, Tarija). However, the Chacoan and Cerrado affinities of the Beni and E Santa Cruz herpetofauna make this assumption too feeble. The taxonomy of the pair of species *L. chaquensis*-*L. macrosternum* is chaotic and needs revision urgently.

Leptodactylus didymus Heyer, García-López & Cardoso, 1996

Distribution: 1) Southwestern Amazonia; 2) Beni, La Paz, **Pando**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: KÖHLER & LÖTTERS, 1999a.

Remarks: *Leptodactylus didymus* and *L. mystaceus* are sibling species barely distinguishable based on morphology. However, the calls of *L. didymus* are more similar to those of *L. elenae* than to those of *L. mystaceus* (MÁRQUEZ *et al.*, 1995; HEYER *et al.*, 1996).

Leptodactylus elenae Heyer, 1978

Distribution: 1) Cerrado, Chaco; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993a; KÖHLER, 1995a; MÁRQUEZ *et al.*, 1995.

Leptodactylus fuscus (Schneider, 1799)

Distribution: 1) Middle and South America; 2) **Beni, Cochabamba, Chuquisaca, La Paz, Pando, Santa Cruz, Tarija;** 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: REYNOLDS & FOSTER, 1992; KÖHLER, 1995a; MÁRQUEZ *et al.*, 1995; KÖHLER & BÖHME, 1996; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Leptodactylus gracilis Duméril & Bibron, 1841

Distribution: 1) Subtropical South America; 2) Cochabamba, **Chuquisaca, Santa Cruz, Tarija;** 3) Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests; 4) Terrestrial; 5) Open.

Key literature: KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; KÖHLER & LÖTTERS, 1999b.

Leptodactylus griseigularis (Henle, 1981)

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz;** 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: HEYER, 1994; KÖHLER, 2000a. Remarks: HEYER (1994) split the former, widespread *L. wagneri* (Peters) into several species. As a result, *L. wagneri* no longer occurs in Bolivia. *Leptodactylus griseigularis* is the name applied to the populations of the Andean slopes and adjacent foothills, whereas *L. leptodactyloides* is the species widespread in the lowlands. Both species are mostly parapatric, but come into contact at the foothills of the Andes.

Leptodactylus knudseni Heyer, 1972

Distribution: 1) Upper Amazon Basin; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz;** 3) Amazonian Rainforests, Humid Transi-

tion Lowland Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1990a.

Leptodactylus labyrinthicus (Spix, 1824)

Distribution: 1) Cerrado; 2) **Beni, Cochabamba, Chuquisaca, Santa Cruz;** 3) Cerrado, Chiquitanía Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993a; KÖHLER, 1995a; 2000a; MÁRQUEZ *et al.*, 1995; KÖHLER & BÖHME, 1996.

Leptodactylus laticeps Boulenger, 1918

Distribution: 1) Chaco; 2) Chuquisaca, Santa Cruz, Tarija; 3) Chaco Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a.

Remarks: FROST (1985) reported the species in Bolivia, but there is no published locality hitherto.

Leptodactylus latinatus Jiménez de la Espada, 1875

Distribution: 1) Chaco; 2) Chuquisaca, **Santa Cruz, Tarija;** 3) Chaco Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a.

Remarks: GALLARDO (1964b) described the subspecies *L. l. anceps*, which would be the one present in Bolivia; this subspecies was invalidated by KLAPPENBACH & LANGONE (1992).

Leptodactylus leptodactyloides (Andersson, 1945)

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz;** 3) Amazonian Rainforests, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; HEYER, 1994; KÖHLER, 1995a; 1995b; 2000a; MÁRQUEZ *et al.*, 1995; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Remarks: Most records compiled by DE LA RIVA (1990a) as *L. wagneri* correspond to *L. leptodactyloides*. See remarks under *L. griseigularis*.

Leptodactylus macrosternum Miranda-Ribeiro, 1926

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA & MALDONADO, 1999.

Remarks: See remarks under *L. chaquensis*.

Leptodactylus mystaceus (Spix, 1824)

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest, Open.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1995.

Leptodactylus mystacinus (Burmeister, 1861)

Distribution: 1) Cerrado, Chaco, Subtropical South America; 2) Chuquisaca, **Santa Cruz**, Tarija; 3) Cerrado, Chaco Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a.

Leptodactylus ocellatus (Linnaeus, 1758)

Distribution: 1) Subtropical South America; 2) **Santa Cruz**; 3) Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA & MALDONADO, 1999.

Remarks: Many old Bolivian records actually corresponding to the pair *L. chaquensis*-*L. macrosternum* were assigned to this species in the past. Its distribution in the country is discussed by DE LA RIVA & MALDONADO (1999).

Leptodactylus pentadactylus (Laurenti, 1768)

Distribution: 1) Tropical Middle and South

America; 2) Beni, Cochabamba, **La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Key literature: APARICIO, 1992.

Leptodactylus petersii (Steindachner, 1864)

Distribution: 1) Amazonia, Cerrado; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: HEYER, 1994; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Remarks: This species was formerly considered a subspecies of *L. podicipinus*. Some of the locality records provided by DE LA RIVA (1990a) for *L. podicipinus* might correspond to *L. petersii*.

Leptodactylus podicipinus (Cope, 1862)

Distribution: 1) Amazonia, Cerrado; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Open.

Key literature: HEYER, 1994; KÖHLER, 1995a; MÁRQUEZ *et al.*, 1995.

Remarks: See remarks under *P. petersii*.

Leptodactylus rhodomystax Boulenger, 1884

Distribution: 1) Upper Amazon Basin; 2) **Beni, La Paz, Pando**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Terrestrial; 5) Forest.

Key literature: FUGLER, 1983.

Leptodactylus rhodonotus (Günther, 1869)

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; KÖHLER, 1995a; 2000a; KÖHLER & LÖTTERS, 1999b.

Leptodactylus syphax Bokermann, 1969

Distribution: 1) Cerrado; 2) **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Open.

Key literature: KÖHLER, 1995a; 1995b; KÖHLER & BÖHME, 1996.

Lithodytes lineatus (Schneider, 1799)

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1993a; MÁRQUEZ *et al.*, 1995.

Physalaemus albonotatus (Steindachner, 1864)

Distribution: 1) Cerrado, Chaco; 2) Beni, **Chuquisaca, Santa Cruz, Tarija**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993a; KÖHLER, 1995a; 2000a; MÁRQUEZ *et al.*, 1995.

Remarks: Many old records of this species were given as *P. cuvieri*, which is very similar (they are indistinguishable in alcohol). See remarks under *P. cuqui*.

Physalaemus biligonigerus (Cope, 1861
“1860”)

Distribution: 1) Cerrado, Chaco; 2) **Chuquisaca, Santa Cruz, Tarija**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests; 4) Terrestrial; 5) Open.

Key literature: MÁRQUEZ *et al.*, 1995; KÖHLER, 2000a.

Remarks: This species has been mistaken for *P. fuscomaculatus* (Steindachner) in the literature for a long period. As a result, the geographic distribution of both species is confused. These problems were discussed by CEI (1990), who clearly stated that these

two names represent different species. PERACCA (1897) reported specimens of both species (as *Paludicola fuscomaculata* Steindachner and *Paludicola signifera* Girard) at some localities in Tarija, but their assignment to one or the other species was not convincing (CEI, 1990). The locality of the rediscovered Paraguayan sample studied by CEI (1990) and, especially, the type locality of *P. fuscomaculatus* (“Caissará”, Mato Grosso do Sul, Brazil) are very close to the Bolivian border. Thus, the presence of the species in the country is very plausible but, according to CEI (1990), *P. fuscomaculatus* is less abundant than *P. biligonigerus* and has a distribution restricted to the lower Paraguay basin. For this reason, the records from Tarija are doubtful and we refrain from including *P. fuscomaculatus* in the Bolivian fauna until conclusive evidence of its occurrence in the country is available.

Physalaemus cuqui Lobo, 1993

Distribution: 1) Chaco; 2) Chuquisaca, **Santa Cruz, Tarija**; 3) Chaco Lowland Forests, Chaco Montane Forests; 4) Terrestrial; 5) Open.

Key literature: LAVILLA & SCROCCHI (1999).

Remarks: This species, very similar to *P. albonotatus*, has been reported in the country only recently. Some Chacoan records of *P. albonotatus* might actually correspond to *P. cuqui*.

Physalaemus cuvieri Fitzinger, 1826

Distribution: 1) Cerrado; 2) **Santa Cruz**; 3) Cerrado; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a.

Remarks: See remarks on *P. albonotatus*.

Physalaemus nattereri (Steindachner, 1863)

Distribution: 1) Cerrado; 2) **Santa Cruz**; 3) Cerrado; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a; KÖHLER, 1995a; MÁRQUEZ *et al.*, 1995.

Physalaemus petersi (Jiménez de la Espada, 1872)

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Santa Cruz**, Pando; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Terrestrial; 5) Forest.

Key literature: CANNATELLA & DUELLMAN, 1984; MÁRQUEZ *et al.*, 1995.

Remarks: CANNATELLA & DUELLMAN (1984) placed *Eupemphix freibergi* Donoso-Barros [type locality: Runerrabaque (sic), Beni] in the synonymy of *Physalaemus petersi*, and in this were followed by subsequent authors. Recently, CANNATELLA *et al.* (1998), in a paper dealing with the phylogeny of the entire *P. pustulosus* species group, again used the name *Physalaemus freibergi*, the distribution of which would be the SW Amazon in Peru and Bolivia. We prefer to maintain the name *P. petersi* for these populations until further studies confirm their taxonomic position.

Pleurodema borellii (Peracca, 1895)

Distribution: 1) Chaco, Temperate Inter-Andean Valleys; 2) Chuquisaca, **Tarija**; 3) Chaco Montane Forests, Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests; 4) Terrestrial; 5) Open.

Key literature: PERACCA, 1897; DE LA RIVA & GONZÁLES, 1998.

Remarks: This species has been considered either as a good species or as a synonym of *P. cinereum*, primarily on the basis of their almost identical morphology. However, it seems odd that a species may range from the hot, lowland Chaco to the high, cold puna (that is, from 400 to 4500 m or more), an argument sometimes utilized to justify the recognition of the two species (DUELLMAN & VELOSO, 1977). However, the biochemical and biacoustical evidence seems to indicate that only one species might be involved (MCMASTER *et al.*, 1991). Argentinean herpetologists tend to recognize two separate species, and populations assigned

to *P. borellii* have been recently found far away from the Andean foothills (CÉSPEDEZ, 1997). Admittedly, less evidence is available in Bolivia to either recognize or reject *P. borellii* as a valid species. Since PERACCA (1897) reported *P. borellii* in Tarija, we tentatively recognize it as valid and include it in the list.

Pleurodema cinereum Cope, 1877

Distribution: 1) Andean Highlands, Temperate Inter-Andean Valleys; 2) **Cochabamba, Chuquisaca, La Paz, Oruro, Potosí**; 3) High Andean Vegetation, Inter-Andean Valleys; 4) Terrestrial; 5) Open.

Key literature: ERGUETA, 1991; VELLARD, 1992; APARICIO, 1993; CHRISTMANN, 1995; DE LA RIVA *et al.*, 1995; IBISCH & BÖHME, 1995; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; MÁRQUEZ *et al.*, 1995; DE LA RIVA & GONZÁLES, 1998.

Remarks: See remarks under *P. borellii*.

Pleurodema guayapae Barrio, 1964

Distribution: 1) Chaco; 2) Chuquisaca, **Santa Cruz**, Tarija; 3) Chaco Lowland Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA & GONZÁLES, 1998; GONZÁLES, 1998.

Remarks: Whereas the presence of this species in poorly surveyed areas of Chuquisaca and Tarija is plausible, the Chacoan regions of northern Argentina are well studied and the occurrence of the species there is discarded. The perplexing distribution of this species, with a gap of 1000 km between the Bolivian and Argentinean populations (which live in salty, arid environments), suggests that they might represent different species, although the morphological evidence does not support such an assumption (DE LA RIVA & GONZÁLES, 1998).

Pleurodema marmoratum (Duméril & Bibron, 1841)*

Distribution: 1) Andean Highlands; 2) Co-

chabamba, La Paz, Oruro, Potosí; 3) High Andean Vegetation; 4) Terrestrial; 5) Open. Key literature: ERGUETA, 1991; REYNOLDS & FOSTER, 1992; VELLARD, 1992; APARICIO, 1993.

Pseudopaludicola boliviana Parker, 1927*
Distribution: 1) Tropical South America; 2) **Beni**, Cochabamba, **La Paz, Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Terrestrial; 5) Open.

Key literature: KÖHLER, 1995a; MÁRQUEZ *et al.*, 1995; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Remarks: LYNCH (1989a) reviewed the *Pseudopaludicola* of N South America, and he could not find differences between specimens from eastern Colombia, southern Venezuela, and Bolivia. The currently accepted distribution of *P. boliviana* comprises eastern Colombia, S Venezuela, Guyana, Surinam, Roraima (N Brazil), E Bolivia, Paraguay, SW Brazil and N Argentina (LYNCH, 1989a; LOBO, 1992; 1994). Thus, there are two sets of populations separated by the Amazon Basin. This strange distribution roughly parallels that of *Phyllomedusa hypochondrialis*. Although it is not necessarily untenable that both sets of populations of *Pseudopaludicola boliviana* are conspecific, further research is necessary to ascertain their phylogenetic relationships and taxonomic status, and perhaps the northern populations might deserve to be considered a different species.

Pseudopaludicola mystacalis (Cope, 1887)
Distribution: 1) Cerrado; 2) **Beni, Santa Cruz**; 3) Cerrado, Wet Savannas; 4) Terrestrial; 5) Open.
Key literature: DE LA RIVA, 1990a.

Vanzolinius discodactylus (Boulenger, 1883)
Distribution: 1) Upper Amazon Basin; 2) Beni,

La Paz, **Pando**; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Key literature: KÖHLER & LÖTTERS, 1999a.

Telmatobiinae

Eleutherodactylus ashkapara Köhler, 2000**
Distribution: 1) Humid Montane Forests; 2) **Cochabamba**, La Paz, Santa Cruz; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.
Key literature: KÖHLER, 2000a; 2000b.

Eleutherodactylus cruralis (Boulenger, 1902)*
Distribution: 1) Bolivian-Argentinean Forests, Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba**, Chuquisaca, **La Paz, Santa Cruz**; 3) Amazonian Rainforests, Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: LYNCH, 1989b; DE LA RIVA, 1993c; MÁRQUEZ *et al.*, 1995; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995b; REICHLE & KÖHLER, 1997; REICHLE *et al.*, 2001.

Remarks: LYNCH (1989b) stated that there is remarkable variation in the size of males and females across the range of the species, a fact that might indicate that more than one species is involved under the name *E. cruralis*. LAVILLA & SCROCCHI (1999) report two unidentified species of the *E. discoidalis* group in Tarija. In our opinion, the morphological variation in samples identified as *E. cruralis* from lowland rainforests in SW Peru and central Bolivia, as well as in those from cloud forests and temperate valleys, are incompatible with the idea of a single species. Bioacoustical evidence also supports the presence of more than one species. A new species is being described by REICHLE *et al.* (2001) and three others will be described by M. Harvey and R. Aguayo.

Eleutherodactylus danae Duellman, 1978
Distribution: 1) Humid Montane Forests, Peri-

Andean Forests; 2) **Cochabamba**, La Paz, Santa Cruz; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: KÖHLER, 1995a; 2000a; KÖHLER & JUNGFER, 1995.

***Eleutherodactylus discoidalis* (Peracca, 1895)**

Distribution: 1) Bolivian-Argentinean Forests, Humid Montane Forests; 2) **Cochabamba, Chuquisaca, Santa Cruz, Tarija**; 3) Tucumanian-Bolivian Montane Forests, Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993c; HARVEY, 1997.

Remarks: DE LA RIVA (1990a; 1993c) overlooked the old record of this species in Tarija by PERACCA (1897). See remarks under *E. cruralis*.

***Eleutherodactylus fenestratus* (Steindachner, 1864)**

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; DE LA RIVA, 1993c; MÁRQUEZ *et al.*, 1995; REICHLE, 1999; KÖHLER, 2000a.

Remarks: Many locality records assigned to this species are probably in error. Considerable confusion exists regarding the taxonomic status of some Peruvian and Bolivian populations of frogs in the *E. conspicillatus* group, which have similar morphologies, moderate to broad distributions, and poorly known regional variation. After REICHLE (1999) *Hylodes gollmeri bisignatus* Werner [type locality: Chaco, Bolivia (a cloud forest locality in Departamento La Paz)], currently in the synonymy of *E. fenestratus*, is a different species whose taxonomic position is still under study (see KÖHLER, 2000a).

Eleutherodactylus fraudator* Lynch & McDiarmid, 1987*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: LYNCH & MCDIARMID, 1987; DE LA RIVA, 1993c; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995a; DE LA RIVA & LYNCH, 1997.

Eleutherodactylus llojsintuta* Köhler & Löters, 1999*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: KÖHLER & LÖTTERS, 1999c; KÖHLER, 2000a.

Remarks: See remarks under *E. platydactylus*.

Eleutherodactylus mercedesae* Lynch & McDiarmid, 1987*

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: LYNCH & MCDIARMID, 1987; DE LA RIVA, 1993c; KÖHLER, 2000a.

Eleutherodactylus olivaceus* Köhler, Morales, Löters, Reichle & Aparicio, 1998

Distribution: 1) Humid Montane Forests, Peri-Andean Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993c; KÖHLER *et al.*, 1998b; KÖHLER, 2000a.

Remarks: This recently described species includes the populations on which former Bolivian records of *E. mendax* Duellman were based (see DE LA RIVA, 1993c). However, the presence of *E. mendax* in the country is still plausible.

Eleutherodactylus peruvianus (Melin, 1941)

Distribution: 1) Humid Montane Forests, Peri-Andean Forests, Upper Amazon Basin; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest. Key literature: DE LA RIVA, 1994b; KÖHLER, 1995a; 2000a.

Remarks: Unlike most Peruvian specimens of *E. peruvianus*, Bolivian ones often lack orange spots on the posterior surfaces of the thighs. There is a considerable variation in these frogs and, given the fact that the type locality of the species (Roque, Department of San Martín, Peru) is far away from Bolivia, it would not be surprising that some Bolivian populations represent a different species (see KÖHLER, 2000a).

Eleutherodactylus platydactylus (Boulenger, 1903)

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993c; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995a; MÁRQUEZ *et al.*, 1995; DE LA RIVA, 1997; KÖHLER & LÖTTERS, 1999c.

Remarks: Recently, DE LA RIVA (1997) placed *E. bockermannii* Donoso-Barros [type locality: Runerrabaque (sic), Beni (probably in error)] in the synonymy of *E. platydactylus*. The striking variation of this species has been pointed out and illustrated repeatedly, and it has been suggested that more than one species might be hidden under the name *E. platydactylus*. DE LA RIVA (1997) refrained from naming new species but described three morphs which might prove to be valid species. *Eleutherodactylus llojsintuta* is a sibling species described by KÖHLER & LÖTTERS (1999c) based mainly on differences in the advertisement calls.

Eleutherodactylus pluvicanorus De la Riva & Lynch, 1997**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995a; DE LA RIVA & LYNCH, 1997.

Eleutherodactylus rhabdolaemus Duellman, 1978

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, La Paz, Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: LYNCH & McDIARMID, 1987; REYNOLDS & FOSTER, 1992; DE LA RIVA, 1993c; KÖHLER, 1995a; 2000a.

Eleutherodactylus samaipatae Köhler & Jungfer, 1995**

Distribution: 1) Bolivian-Argentinean Forests, Temperate Inter-Andean Valleys; 2) Cochabamba, Chuquisaca, **Santa Cruz, Tarija**; 3) Inter-Andean Valleys, Tucumanian-Bolivian Montane Forests; 4) Terrestrial, Arboreal; 5) Forest, Open.

Key literature: KÖHLER, 1995a; 2000a; KÖHLER & JUNGFER, 1995; KÖHLER *et al.*, 1995b; REICHLE, 1999.

Eleutherodactylus toftae Duellman, 1978

Distribution: 1) Peri-Andean Forests; 2) **Cochabamba**, La Paz, Santa Cruz; 3) Amazonian Rainforests; 4) Terrestrial, Arboreal; 5) Forest.

Key literature: DE LA RIVA, 1993c; MÁRQUEZ *et al.*, 1995.

Eleutherodactylus ventrimarmoratus (Boulenger, 1912)

Distribution: 1) Humid Montane Forests, Upper Amazon Basin; 2) Beni, La Paz, **Pando**; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal; 5) Forest.

Remarks: Reported herein for the first time for Bolivia.

Eleutherodactylus zongoensis Reichle & Köhler, 1997**

Distribution: 1) Humid Montane Forests; 2) **La Paz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REICHLE & KÖHLER, 1997.

Remarks: Known only from the holotype.

Ischnocnema quixensis (Jiménez de la Espada, 1872)

Distribution: 1) Amazonia; 2) Beni, La Paz, **Pando**; 3) Amazonian Rainforests; 4) Terrestrial; 5) Forest.

Remarks: Found by J. E. Cadle and S. Reichle in November 1999 and reported herein for the first time.

Ischnocnema sanctaecrucis Harvey & Keck, 1995**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba**, La Paz, **Santa Cruz**; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: HARVEY & KECK, 1995; REICHLE, 1999; KÖHLER, 2000a.

Odontophrynus americanus (Duméril & Bibron, 1841)

Distribution: 1) Subtropical South America; 2) **Cochabamba**, **Chuquisaca**, **Santa Cruz**, **Tarija**; 3) Chaco Lowland Forests, Chaco Montane Forests; 4) Fossorial, Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a; MÁRQUEZ *et al.*, 1995; KÖHLER, 2000a.

Odontophrynus lavillai Cei, 1985

Distribution: 1) Chaco, Temperate Inter-Andean Valleys; 2) Chuquisaca, **Santa Cruz**, Tarija; 3) Chaco Lowland Forests, Inter-Andean Valleys; 4) Fossorial, Terrestrial; 5) Open.

Key literature: DE LA RIVA *et al.*, 1996c; KÖHLER, 2000a.

Remarks: The distribution and variation of *O. americanus* and *O. lavillai* in the country is

still unclear, and both forms seem to be sympatric in some areas.

Phrynoporus kempffi De la Riva, 1992**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba**, **Santa Cruz**; 3) Ceja de Montaña-Cloud Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1992b; MÁRQUEZ *et al.*, 1995.

Phrynoporus laplacai (Cei, 1968)**

Distribution: 1) Humid Montane Forests; 2) **La Paz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest, Open.

Key literature: CEI, 1968; LYNCH, 1975; ERGUETA, 1993.

Phrynoporus pinguis Harvey & Ergueta, 1998**

Distribution: 1) Humid Montane Forests; 2) **La Paz**; 3) Ceja de Montaña-Cloud Forests; 4) Terrestrial; 5) Forest.

Key literature: HARVEY & ERGUETA, 1998.

Remarks: The genus *Phrynoporus* seems to be much more speciose in Bolivia than previously expected. Two new species are being described by R. Aguayo and M. Harvey, and some more will be described by I. De la Riva and S. Reichle.

Phyllonastes carrascoicola De la Riva & Köhler, 1998**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba**, **Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: REYNOLDS & FOSTER, 1992; KÖHLER, 1995a; 2000a; 2000c; KÖHLER *et al.*, 1995a; DE LA RIVA & KÖHLER, 1998.

Phyllonastes ritarasquinae Köhler, 2000**

Distribution: 1) Humid Montane Forests; 2) **Cochabamba**, La Paz, Santa Cruz; 3) Yungas-Montane Rainforests; 4) Terrestrial; 5) Forest.

Key literature: KÖHLER, 2000a; 2000c.

Remarks: More species of the genus *Phyllostomastes* [or a closely related minute leptodactylid genus (M. B. Harvey, in litt.)] are expected to be found in the country as closer attention be paid to the forest litter amphibian fauna.

Telmatobius boliviensis* Parker, 1940*

Distribution: 1) Humid Montane Forests; 2) **La Paz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic; 5) Forest.

Key literature: PARKER, 1940; VELLARD, 1951; 1970; LAVILLA & DE LA RIVA, 1993.

Remarks: The taxonomy of the Yungas *Telmatobius* is obscured by historical problems related to unclear type localities, state of flux in relation to synonymizations and revalidations, the intrinsic difficulty posed by the frogs themselves due to their intra- and interpopulational variation, and the scarcity of museum specimens. *Telmatobius boliviensis* was placed in the synonymy of *T. verrucosus* Werner by VELLARD (1951), who later considered both forms as subspecies of *T. marmoratus* (VELLARD, 1970). DE LA RIVA (1990a) considered doubtful the validity of *T. verrucosus* (and, as a consequence, that of *T. boliviensis*). Later, LAVILLA & DE LA RIVA (1993) described the putative tadpole of *T. boliviensis*. The tadpole was assigned to this species on the basis of the comparison of some adult frog specimens (for which the tadpole was known) with the holotype of *T. boliviensis*. Further observations led to conclude that more than one species is involved and the tadpoles were wrongly assigned (I. De la Riva, pers. obs.). *Telmatobius boliviensis* and *T. verrucosus* seem to be different species (but see remarks under *T. verrucosus*) and at least there is another related species, which is being described by I. De la Riva.

Telmatobius culeus* (Garman, 1875)

Distribution: 1) Andean Highlands; 2) **La**

Paz; 3) High Andean Vegetation; 4) Aquatic; 5) Open.

Key literature: BARBOUR & NOBLE, 1920; PARKER, 1940; VELLARD, 1951; 1953; 1955; 1960; DE MACEDO, 1960; CEI, 1986; VELLARD, 1992; PÉREZ, 1998.

Remarks: The description of subspecies on the basis of morphological differences has created a taxonomic chaos in some species groups of *Telmatobius*. Up to six subspecies of *T. culeus* have been recognized. The nominal subspecies has its type locality in Bolivia [Achacache, Lago Titicaca (La Paz) (FROST, 1985) or "Proximidades de la Isla del Sol" (La Paz) (VELLARD, 1953)]. Furthermore, VELLARD (1951; 1960) described three subspecies of *T. albiventris* Parker, a species that he later placed in the synonymy of *T. culeus*, but he still recognized all the former subspecies of *T. albiventris* (VELLARD, 1992). One of these subspecies, *T. a. globulosus* Vellard, has its type locality in Bolivia [Playa de Copani, Huiñamarca (La Paz)]. As a result, the number of subspecies of *T. culeus* rose to twelve. Fortunately, a thorough study carried out by SINSCH *et al.* (1995) demonstrated that the morphological variation in *Telmatobius* does not match subspecific partitions like this. We follow VELLARD (1992) in that *T. albiventris* is a synonym of *T. culeus*.

***Telmatobius edaphonastes* De la Riva, 1995
"1994" ****

Distribution: 1) Humid Montane Forests; 2) **Cochabamba, Santa Cruz**; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic, Terrestrial; 5) Forest.

Key literature: DE LA RIVA (1995c).

Telmatobius huayra* Lavilla & Ergueta, 1995*

Distribution: 1) Andean Highlands; 2) **Potosí**; 3) High Andean Vegetation; 4) Aquatic; 5) Open.

Key literature: LAVILLA & ERGUETA, 1995a.

Telmatobius ifornoi Lavilla & Ergueta, 1999**

Distribution: 1) Humid Montane Forests; 2)

La Paz; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic; 5) Forest.

Key literature: LAVILLA & ERGUETA, 1999.

Telmatobius jahuiira Lavilla & Ergueta, 1995**

Distribution: 1) Humid Montane Forests; 2)

La Paz; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic, Terrestrial; 5) Forest.

Key literature: LAVILLA & ERGUETA, 1995b.

Telmatobius marmoratus (Duméril & Bibron, 1841)

Distribution: 1) Andean Highlands; 2) **Cochabamba, Chuquisaca, La Paz, Oruro, Potosí;** 3) High Andean Vegetation, Inter-Andean Valleys; 4) Aquatic; 5) Open.

Key literature: PARKER, 1940; VELLARD, 1951; 1953; 1955; 1960a; 1969; 1970; 1992; CEI, 1986; ERGUETA, 1991; APARICIO, 1993; IBISCH & BÖHME, 1993; KÖHLER, 1995a.

Remarks: This is the most widespread species of *Telmatobius* and the one with the most complicated taxonomy. Once *T. boliviensis* is resurrected as a valid species, still nine subspecies of *T. marmoratus* remain. Two of them have their type locality in Bolivia: *T. m. hintoni* Parker (Tolota, Cochabamba) and *T. m. gigas* Vellard (Huayllamarca, Oruro). Furthermore, the nominal subspecies and *T. m. rugosus* Vellard would also occur in the country. The validity of all these subspecies and the real specific diversity of the *T. marmoratus* complex in Bolivia is being reviewed by I. De la Riva.

Telmatobius simonsi Parker, 1940**

Distribution: 1) Temperate Inter-Andean Valleys; 2) **Cochabamba, Chuquisaca, Potosí, Santa Cruz,** Tarija; 3) Inter-Andean Valleys; 4) Aquatic; 5) Open.

Key literature: PARKER, 1940; VELLARD, 1951.

Remarks: Certain populations of cloud forest *Telmatobius* in Cochabamba and Santa Cruz were tentatively identified as *T. simonsi* (DE LA RIVA, 1990a; 1994c). These populations actually represent an unnamed species, which is being described by I. De la Riva and M. Harvey, who are also reviewing the taxonomic status of some other populations allied to *T. simonsi* from the temperate valleys of Santa Cruz.

Telmatobius verrucosus Werner, 1899**

Distribution: 1) Humid Montane Forests; 2)

La Paz; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic; 5) Forest.

Key literature: WERNER, 1899; BARBOUR & NOBLE, 1920; VELLARD, 1951; 1970.

Remarks: This species has not been found again, and its uncertain taxonomic status is being investigated by I. De la Riva. See remarks under *T. boliviensis*.

Telmatobius yuracare De la Riva, 1994**

Distribution: 1) Humid Montane Forests; 2)

Cochabamba, Santa Cruz; 3) Ceja de Montaña-Cloud Forests, Yungas-Montane Rainforests; 4) Aquatic; 5) Forest.

Key literature: DE LA RIVA, 1994c; KÖHLER, 1995a; 2000a; KÖHLER *et al.*, 1995a.

Microhylidae**Microhylinae*****Chiasmocleis albopunctata*** (Boettger, 1885)

Distribution: 1) Cerrado; 2) **Santa Cruz;** 3)

Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1996c; KÖHLER, 2000a.

Chiasmocleis ventrimaculata (Andersson, 1945)

Distribution: 1) Upper Amazon Basin; 2) **Beni, Cochabamba**, La Paz, Pando, Santa Cruz; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1995d.

Remarks: WALKER (1973) reported a specimen of *C. bassleri* Dunn from Bolivia. So far, *C. bassleri* is known from E Peru and Ecuador (FROST, 1985), but new findings in Brazil, close to the Bolivian border, make Walker's report plausible (S. Reichle, pers. obs.).

Dermatonotus muelleri (Boettger, 1855)

Distribution: 1) Chaco, Cerrado; 2) Chuquisaca, **Santa Cruz, Tarija**; 3) Chaco Lowland Forests; 4) Fossal, Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1990a.

Elachistocleis bicolor (Valenciennes, 1838)

Distribution: 1) Cerrado, Chaco, Southwestern Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Fossal, Terrestrial; 5) Open.

Key literature: KÖHLER, 1995a; 2000a; 1995b; DE LA RIVA *et al.*, 1996c; REICHLE, 1996; 1997c; REICHLE & KÖHLER, 1998.

Remarks: Considerable confusion took place in the past regarding the identity of *E. bicolor* vs. *E. ovalis*, whose main external difference is the ventral coloration (see FROST, 1985). Today most authors apply the name *E. bicolor* to the frogs with an immaculate venter.

Elachistocleis ovalis (Schneider, 1799)

Distribution: 1) South America; 2) **Beni, Cochabamba, Chuquisaca, La Paz, Pando**,

Santa Cruz; 3) Amazonian Rainforests, Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Chaco Montane Forests, Humid Transition Lowland Forests, Inter-An- dean Valleys, Pre-Cambrian Shield Humid Forests, Tucumanian-Bolivian Montane Forests, Wet Savannas; 4) Fossal, Terrestrial; 5) Open.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1996c; REICHLE, 1997c; KÖHLER, 2000a.

Remarks: See comments under *E. bicolor*.

Hamptophryne boliviensis (Parker, 1927)*

Distribution: 1) Upper Amazon Basin; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1993a; DE LA RIVA *et al.*, 1996c.

Pipidae

Pipinae

Pipa pipa (Linnaeus, 1758)

Distribution: 1) Amazonia; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Aquatic; 5) Forest, Open.

Key literature: REICHLE, 1997a; HARVEY *et al.*, 1998.

Pseudidae

Lysapsus limellus Cope, 1862

Distribution: 1) Cerrado, Lower Amazon Basin; 2) **Beni, Cochabamba, La Paz, Santa Cruz, Pando**; 3) Cerrado, Wet Savannas; 4) Aquatic; 5) Open.

Key literature: KÖHLER, 1995a; BOSCH *et al.*, 1996; REICHLE, 1996; 1997a; REICHLE & KÖHLER, 1998.

Remarks: The taxonomy of *L. limellus* was reviewed by GALLARDO (1961b; 1964c) but the status of several populations remains problematic. In Bolivia the subspecies *L. limellus boliviensis* Gallardo (type locality: Reyes, Beni) occurs, and probably *L. l. limellus*, whose type locality (Corumbá, Mato Grosso do Sul) lies on the Brazilian bank of the Paraguay river, which divides the Brazilian and Bolivian Pantanal. KLAPPENBACH (1985) stated that *L. l. laevis* Parker, from the lower Amazon, should be considered a good species, and suggested that *L. l. boliviensis* might be either a valid species or a subspecies of *L. laevis*. At least two different morphs corresponding to two kinds of advertisement calls are found in the Beni area (S. REICHLE, pers. obs.).

Pseudis paradoxa (Linnaeus, 1758)

Distribution: 1) South America; 2) **Beni**, Cochabamba, Chuquisaca, La Paz, Pando, **Santa Cruz**; 3) Cerrado, Chiquitanía Forests, Chaco Lowland Forests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests, Wet Savannas; 4) Aquatic; 5) Open.

Key literature: BOSCH *et al.*, 1996; REICHLE, 1996; 1997a; REICHLE & KÖHLER, 1998; DE LA RIVA, 1999d.

Remarks: Three subspecies have been reported in the country: *P. p. paradoxa* (Linnaeus) in the Beni region, *P. p. platensis* Gallardo in the Cerrado and *P. p. occidentalis* Gallardo (type locality: El Pailón, Santa Cruz) in the Chaco (GALLARDO, 1961b; 1964c). As stated by BOSCH *et al.* (1996) and DE LA RIVA (1999d), the validity and distribution of these forms in the country needs to be studied, especially the presence of the nominal subspecies, whose core area of distribution is the Guianan region and for which there are no records in the intervening Amazonian region.

Ranidae

Raninae

Rana palmipes Spix, 1824

Distribution: 1) Amazonia; 2) **Beni**, **Cochabamba**, **La Paz**, Pando, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Terrestrial; 5) Forest.

Key literature: DE LA RIVA, 1990a.

Caudata

Plethodontidae

Plethodontinae

Bolitoglossa altamazonica (Cope, 1874)

Distribution: 1) Amazonia; 2) Beni, **Cochabamba**, La Paz, **Santa Cruz**, Pando; 3) Amazonian Rainforests, Yungas-Montane Rainforests; 4) Arboreal, Terrestrial; 5) Forest.

Key literature: REICHLE *et al.*, 2000.

Remarks: *Bolitoglossa altamazonica* is thought to include populations from the mouth of the Amazon to the Andean cloud forests from Colombia to Bolivia (there are scant records from the central Amazon). This is one of the least known species complex of South American amphibians, and it is expected that, upon future research, the Bolivian populations will be identified as one or more different species (REICHLE *et al.*, 2000).

Gymnophiona

Caeciliidae

Caecilia marcusi Wake, 1984**

Distribution: 1) Peri-Andean Forests, Southwestern Amazonia; 2) Beni, **Cochabamba**, La Paz, **Santa Cruz**; 3) Amazonian Rainforests, Humid Transition Lowland Forests; 4) Fossorial; 5) Forest.

Key literature: WAKE, 1984; REICHLE & KÖHLER, 1996b.

***Siphonops annulatus* (Mikan, 1820)**

Distribution: 1) Tropical South America; 2) **Beni, Cochabamba, La Paz, Pando, Santa Cruz;** 3) Amazonian Rainforests, Humid Transition Lowland Forests, Pre-Cambrian Shield Humid Forests; 4) Fossal; 5) Forest.

Key literature: DE LA RIVA, 1990a.

***Siphonops paulensis* Boettger, 1892**

Distribution: 1) Cerrado; 2) Beni, **Santa Cruz;** 3) Cerrado, Chiquitania Forests, Chaco Lowland Forests; 4) Fossal; 5) Open.

Key literature: DE LA RIVA, 1990a.

CONCLUSIONS AND DISCUSSION

Taxonomic diversity and changes since 1990.

In total, up to date, there are three orders, eleven families, 44 genera and 186 valid species of amphibians recognized from Bolivia. Their taxonomic distribution is summarized in Table 1. These figures represent a dramatic increase with respect to DE LA RIVA's (1990a) list. Frogs belonging to nine genera not previously reported in the country have been found (either as undescribed species or as already known ones): *Dendrophryniscus*, *Melanophryniscus*, *Allobates*, *Scarthyla*, *Chacophrys*, *Lepidobatrachus*, *Vanzolinius*, *Ischnocnema*, and *Phyllonastes*. As could be expected, the most significant change is that of the total number of species, which has increased by 74, from 112 to 186. This change, which overall represents an increase of 66 % in only ten years, is due to several factors, namely: species overlooked in the 1990 list, species deletions, taxonomic changes, rediscovery of doubtful forms that proved to be valid species, species new to science, and new species records for the country.

Species overlooked in the 1990 list.-*Leptodactylus macrosternum*, *Pleurodema borellii*, and *Eleutherodactylus discoidalis*.

Species deletions.-There are some species deleted from the 1990 list but, due to taxonomic or nomenclatural reasons, they have been replaced by another species' name (e. g., *Scinax parkeri* vs. *S. fuscomarginatus*). In other cases, they have been identified as new species and described (e. g., *Eleutherodactylus olivaceus* vs. *E. mendax*). These cases do not alter the number of species. However, there are two cases of species present in the 1990 list that have been completely removed from the present one: the hylids *Hemiphractus johnsoni* (Noble) and *Hyla rossalleni*. The case of *H. rossalleni* has been already commented in the remarks section of *Hyla leali*. There are two references suggesting the presence of *Hemiphractus johnstoni* in Bolivia. BOULENGER (1903a) treated the reproduction of this species (as *Ceratohyla bubalus*) based on a specimen obtained by Mr. Ockenden at Santo Domingo (Puno, Peru) and commented that it is "...an inhabitant of the Andes of Ecuador, Bolivia and Peru". However, as far as we know, no voucher specimens from Bolivia are available. TRUEB (1974) listed a purported Bolivian specimen of *H. johnstoni* from "Linimbane", 1000 m. [housed at the Natural History Museum, London (BM)]. The name of this locality is an obvious misspelling corresponding either to Limbani or, more probably, to the Linimbare (=Inambari) river. Both sites, in the Peruvian department of Puno, were visited in 1900 by P. O. Simons, collector of the BM. These localities, as well as Santo Domingo, are not far away from the Bolivian border (ca.70 km by airline distance). Thus, *H. johnstoni* is likely to occur in the country, but its presence is still to be confirmed.

Taxonomic changes.-Three kinds of taxonomic changes have taken place: (1) synonymizations (the synonymized "species" disappears from the list); (2) change from subspecific to specific status (a species is added if more than one subspecies were already cited); and (3)

Table 1: Taxonomic diversity and number of endemic species of Amphibians of Bolivia.
Tabla 1: Diversidad taxonómica y número de especies endémicas de anfibios de Bolivia.

Order	Family	Genus	Spp	Endemics
Anura	Bufoidae	<i>Atelopus</i>	1	0
		<i>Bufo</i>	14	2
		<i>Dendrophryniscus</i>	1	0
		<i>Melanophryniscus</i>	1	0
			—	—
			4	2
			—	—
Centrolenidae		<i>Cochranella</i>	3	2
		<i>Hyalinobatrachium</i>	1	0
			—	—
			2	2
Dendrobatidae		<i>Allobates</i>	1	0
		<i>Colostethus</i>	3	1
		<i>Epipedobates</i>	4	1
			—	—
			3	2
			—	—
Hylidae		<i>Gastrotheca</i>	4	2
		<i>Hyla</i>	32	3
		<i>Osteocephalus</i>	4	0
		<i>Phrynohyas</i>	3	0
		<i>Scarthyla</i>	1	0
		<i>Scinax</i>	10	0
		<i>Sphaenorhynchus</i>	1	0
		<i>Phyllomedusa</i>	8	0
			—	—
			8	5
			—	—
Leptodactylidae		<i>Ceratophrys</i>	2	0
		<i>Chacophrys</i>	1	0
		<i>Lepidobatrachus</i>	1	0
		<i>Adenomera</i>	3	0
		<i>Leptodactylus</i>	23	0
		<i>Lithodytes</i>	1	0
		<i>Physalaemus</i>	6	0
		<i>Pleurodema</i>	4	0
		<i>Pseudopaludicola</i>	2	0
		<i>Vanzolinia</i>	1	0
		<i>Eleutherodactylus</i>	17	7

Table 1: Continuation.**Tabla 1:** Continuación.

Order	Family	Genus	Spp	Endemics
		<i>Ischnocnema</i>	2	1
		<i>Odontophrynus</i>	2	0
		<i>Phrynobius</i>	3	3
		<i>Phyllonastes</i>	2	2
		<i>Telmatobius</i>	10	8
			—	—
		16	80	21
	Microhylidae	<i>Chiasmocleis</i>	2	0
		<i>Dermatonotus</i>	1	0
		<i>Elachistocleis</i>	2	0
		<i>Hamptophryne</i>	1	0
			—	—
		4	6	0
	Pipidae	<i>Pipa</i>	1	0
	Pseudidae	<i>Lysapsus</i>	1	0
		<i>Pseudis</i>	1	0
			—	—
		2	2	0
	Ranidae	<i>Rana</i>	1	0
Caudata	Plethodontidae	<i>Bolitoglossa</i>	1	0
Gymnophiona	Caeciliidae	<i>Caecilia</i>	1	1
		<i>Siphonops</i>	2	0
			—	—
		2	3	1
3	11	44	186	33

resurrection from synonymy (a species is added if the senior synonym represents a species also present in the country).

Synonymizations affected eight species: *Atelopus willimani*, *Hyla callipleura*, *H. carinata* (DE LA RIVA, 1993d), *H. ocapia*, *H.*

steinbachi (it was listed in the 1990 list, but synonymized with *H. fasciata* therein), *Oolygon* (= *Scinax*) *epacrorthina*, *Eleutherodactylus bockermannii*, and *Telmatobius albiventris*.

Case (2) affects five species: *Bufo poepigii*, *Epipedobates hahneli*, *Hyla andina*,

Leptodactylus petersii, and *Telmatobius boliviensis*.

Case (3) is represented by four species: *Scinax parkeri*, *Adenomera diptyx*, *Leptodactylus griseigularis*, and *L. leptodactyloides*.

Rediscovery of doubtful forms.-Two species listed as doubtful in 1990 are included in the present list: *Gastrotheca splendens* and *Telmatobius verrucosus*.

Species new to science.-A total of 28 Bolivian species of anurans in 13 genera have been described as new since 1990: *Bufo amboroensis*, *B. castaneoticus* (described from Brazil), *B. justinianoi*, *Cochranella nola*, *Gastrotheca lauzuricae*, *Hyla chlorostea*, *H. mariannitae* (described from Argentina), *Scinax castroviejoi*, *S. chiquitanus*, *Phyllomedusa camba*, *Leptodactylus didymus* (described from Peru), *Physalaemus cuqui* (described from Argentina), *Eleutherodactylus ashkapara*, *E. llojsintuta*, *E. olivaceus*, *E. pluvicanorus*, *E. samaipatae*, *E. zongoensis*, *Ischnocnema sanctaecrucis*, *Phrynopus kempffi*, *P. pinguis*, *Phyllonastes carrascoicola*, *P. ritarasquinae*, *Telmatobius edaphonastes*, *T. huayra*, *T. ifornoi*, *T. jahuira*, and *T. yuracare*.

All of them except those described from other countries plus *Bufo amboroensis* (which occurs in Peru), *S. cinax castroviejoi* (which presumably occurs in Argentina), *Phyllomedusa camba* (which occurs in Peru and Brazil), and *Eleutherodactylus olivaceus* (which occurs in Peru) are Bolivian endemisms. The number of endemic species is now 33, which represents 17.7 % of the total amphibian fauna of the country. Most of these endemic species (21) belong to the family Leptodactylidae.

Finally, two species described in the last decade were considered as not valid in the same period: *Bufo echinodes* Reynolds & Foster, and *Centrolenella flavidigitata* Reynolds & Foster were, respectively, synonymized with *B. quechua* and *Cochranella bejaranoi*

by HARVEY & SMITH (1993) and HARVEY (1996).

Overall, of 50 valid species (at present-day) described from Bolivia over a period of 162 years (in 1838 was described the first species whose type locality is in Bolivia, *Epipedobates pictus*), 25 (50%) have been described since 1990. This figure gives an idea about how much the research in Bolivia has lately become dynamic and intense in comparison to past periods. New species are expected to be described in the next years, especially, but not exclusively, in the genera *Bufo*, *Hyla*, *Eleutherodactylus*, *Phrynopus*, and *Telmatobius*.

New species for the country.-They are 37 species in 22 genera, accounting for the largest number of additions to the 1990 list. These species are: *Bufo guttatus*, *Dendrophryniscus minutus*, *Melanophrynniscus rubriventris*, *Cochranella pluvialis*, *Allobates femoralis*, *Colostethus brunneus*, *C. trilineatus*, *Epipedobates hahneli*, *E. trivittatus*, *Hyla albopunctata*, *H. koechlini*, *H. marmorata*, *H. sarayacuensis*, *H. schubarti*, *H. tritaeniata*, *Phrynohyas resinifictrix*, *Scarthyla ostiodactyla*, *Scinax squalirostris*, *Phyllomedusa bicolor*, *P. palliata*, *P. tomopterna*, *Chacophrys pierottii*, *Lepidobatrachus laevis*, *Leptodactylus gracilis*, *L. ocellatus*, *L. pentadactylus*, *L. syphax*, *Physalaemus cuqui*, *Pleurodema guayapae*, *Vanzolinia discodactylus*, *Eleutherodactylus danae*, *E. peruvianus*, *E. platydactylus*, *E. toftae*, *Ischnocnema quixensis*, *Odontophrynuus lavillai*, and *Chiasmocleis ventrimaculata*.

The occurrence of 19 of these species was expected by DE LA RIVA (1990a). The number of new records increases steadily with every herpetological field trip to poorly surveyed areas of the country, especially those close to the border of neighboring countries.

Two species of frogs reported recently in Bolivia have not been included in the present list. *Hyla walfordi* Bokermann was reported in the country by DE LA RIVA *et al.* (1997), based

mainly on bioacoustical criteria. Further examination of the specimens and comparative studies of their advertisement calls revealed that the reported population actually corresponds to *H. tritaeniata* (I. De la Riva, pers. obs.). Thus, *H. walfordi* has not been included in the present list, but its occurrence in the country is highly plausible [see LANGONE & BASSO (1987)]. *Phrynoporus peruvianus* (Noble) was reported by KÖHLER (1995a; b), but further examination of the single specimen collected demonstrated that it was a juvenile of *Pleurodema marmoratum* (I. De la Riva & S. Reichle, pers. obs.).

Species of doubtful status.

For many species named long ago there may be no more published information than that provided in the original description. However, these species may be common and/or have broad distributions. The redescription and publication of biological or ecological data on these species is always advisable. More remarkable are those cases in which a species has remained unknown for a long period and the expectations to get new information are scant, due to the loss of the type material or because their type locality is unknown. We are unhappy with lists containing species whose information is almost untrievable. Such lists obscure the reality of what is well known about the composition and diversity of a given fauna. However, we are aware of the importance of taking *species inquirenda* into account in order to carry out further taxonomic research properly. The case of *Gastrotheca splendens* commented above is an illustrative example: being unaware of the existence of such a "lost" taxon, probably would have led the taxonomist to describe a new species. Thus, a briefly commented list of Bolivian (or putatively Bolivian) doubtful species follows:

Bufo simus Schmidt, 1857. A case equal to that of *Gastrotheca splendens* [see SAVAGE (1972)].

Hyla molitor Schmidt, 1857. A case equal to that of *Gastrotheca splendens* [see DUELLMAN (1970)].

Hyla roeschmanni De Grys, 1938. Type lost and type locality imprecise ("Beni, Bolivia").

Leptodactylus andicola Boettger, 1891. Type lost and generic assignment doubtful (see DE LA RIVA, 1990a).

Species expected to occur.

Many species not yet discovered are expected within the Bolivian boundaries when further surveys are done. Some of the species listed below have been reported very close to the Bolivian border; others are known from more distant localities, but substantial range extensions into Bolivia are plausible. In spite of the fact that many species predicted by DE LA RIVA (1990a) have already been found in Bolivia (see above), the number of potential species (we propose 67) remains roughly as high as in 1990. This is due to two facts: the improvement of the knowledge on the distribution of many species and the description of several new species from localities close to Bolivia. In DE LA RIVA's (1990a) arrangement, there were four different geographical areas in Bolivia to be considered in relation to species potentially occurring in the country. Here we add a fifth zone, the Cordillera Occidental. The areas and potential species are:

Amazonia: *Allophryne ruthveni*, *Bufo glaberrimus*, *Colostethus peruvianus*, *Dendrobates biolat*, *D. quinquevittatus*, *D. vanzolinii*, *Epipedobates macero*, *E. simulans*, *Hyla altenorum*, *H. bokermanni*, *H. brevifrons*, *H. rossalleni*, *H. timbeba*, *H. walfordi*, *H. xapuriensis*, *Scinax ictericus*, *S. pedromedinae*, *Sphaenorhynchus carneus*, *Osteocephalus subtilis*, *Agalychnis craspedopus*, *Phyllomedusa atelopoides*, *Edalorhina perezi*, *Hydrolaetare schmidti*, *Leptodactylus dantasi*, *Eleutherodactylus altamazonicus*, *E. buccinator*, *E. imitatrix*, *E. skydmarios*, *E. zimmermanae*, *Phyllonastes myrmecoides*, *Phyzelaphryne miriamae*, *Altigius alios*, *Chiasmocleis bassleri*, *C.*

shudikarensis, *Ctenophryne geayi*, *Caecilia tentaculata*, *Oscaecilia bassleri*, and *Typhlonectes compressicauda*.

Chaco and Cerrado: *Melanophryniscus stelzneri*, *Epipedobates braccatus*, *Hyla varelae*, *Lepidobatrachus llanensis*, *Physalaemus centralis*, *P. fuscomaculatus*, *Eleutherodactylus dundeei*, and *Chiasmocleis meheleyi*.

Northern Cordillera Oriental: *Atelopus erythropus*, *Cochranella phenax*, *C. spiculata*, *C. truebae*, *Hemiphractus johnsoni*, *Eleutherodactylus cosnippatae*, *E. lindae*, *E. mendax*, *E. ockendeni*, *E. salaputum*, *Phrynoporus peruvianus*, and *Telmatobius sanborni*.

Southern Cordillera Oriental: *Bufo gallardoi*, *B. gnustae*, *B. rumbolli*, *Gastrotheca chrysosticta*, *G. gracilis*, and *Telmatobius oxycephalus*.

Cordillera Occidental: *Telmatobius halli*, *T. pefauri*, and *T. zapahuirensis*.

Conservation, recommendations and future research.

The figure of 186 species of amphibians in Bolivia is still very conservative. Considering the expected descriptions of new species plus the finding of species known from neighboring areas of other countries, a more realistic figure for the Bolivian amphibian diversity is around 250-300 species. The majority of the endemic amphibians of Bolivia occur in the Cordillera Oriental of the Andes, and have restricted distributions in fragile ecosystems (see KÖHLER *et al.*, 1998b; KÖHLER 2000a). The existence of this rich endemic fauna must be necessarily taken into account by Bolivian agencies involved in the management and protection of natural resources, in order to make sound and responsible environmental policy. This amphibian endemism often parallels equally interesting faunas and floras, but many groups of animals and plants remain poorly investigated. Thus, in general, preserving areas of amphibian endemism warrants the preservation of other valuable biological resources. However, the amphibians are often ignored

and mostly excluded from biodiversity inventories, faunal lists, and regional surveys, primarily due to lack of information and taxonomic expertise. For example, The Red Data Book of Bolivian Vertebrates (ERGUETA & MORALES, 1996) includes only two species of amphibians (*Telmatobius albiventris* and *T. culeus*, which are synonyms), and contains recommendations about some others, but these recommendations are only tentative, for sound knowledge of the real problems affecting Bolivian amphibians is lacking. Financial support and academic stimulus to carry out herpetological research are paramount if amphibians are to occupy the place they deserve in conservation policies. Especially, the situation in Bolivia concerning the global phenomenon of declining amphibian populations is completely unknown. Some observations seem to indicate that drastic declines might be affecting certain populations of montane species [e. g., some *Telmatobius*; (ERGUETA & HARVEY, 1996; I. De la Riva, pers. obs.)]. Locally, rivers and lakes are being polluted and large tracts of forest are being wiped out by agriculture, mining, logging, ranching, erosion, road construction, oil exploitation, etc. Information on amphibian diversity and distribution is a first step to know what is being destroyed. Nevertheless, overall, Bolivia is still well preserved and, if natural resources are wisely managed, there are still great opportunities to leave a rich natural heritage to future generations. We hope that further updated lists of Bolivian amphibians will not have to include a section of "probably extinct" species.

Acknowledgments

Understandably, our years of work on Bolivian amphibians which resulted in this list would not have been possible without the kind help of many friends and colleagues. To list all of them would probably require a separate publication. Nevertheless, we want to thank all these persons and institutions who helped

with loaning specimens, providing miscellaneous information, or supporting our work in any other way. We specially thank the kind collaboration and support provided by J. Aparicio and the personnel at the Colección Boliviana de Fauna, La Paz, the Museo de Historia Natural "Noel Kempff Mercado", Santa Cruz, and the Fundación Amigos de la Naturaleza, Santa Cruz. This paper greatly benefitted from the kindness of J. Dorda, W. Duellman, M. Göd, L. Gonzáles, P. Ibisch, K.-H. Jungfer, A. Kwet, the late A. Mújica, H. Nickel, D. Rumiz, and E. Wild, who loaned some slides, and from the patience and efficient management by the REH's editors. The manuscript was improved by the wise comments provided by M. B. Harvey, M. S. Hoogmoed and J. A. Langone. Fieldwork of JK and SL was funded by grants of the German Academic Exchange Service (DAAD) and the "Graduiertenförderung des Landes Nordrhein-Westfalen". This paper is partially funded by project DGES PB97-1147, Spanish Ministry of Education and Culture (I. De la Riva, principal investigator).

REFERENCES

- ANDERSSON, L. G. (1906): On batrachians from Bolivia, Argentina and Peru, collected by Erland Nordenskiöld 1901-1902 and Nils Holmgren 1904-1905. *Ark. Zool.*, 3 (12): 1-20.
- ANDERSSON, L. G. (1938): Batrachians from Burma collected by Dr. R. Malaise and from Bolivia and Ecuador collected by Dr. Hammarlund. *Ark. Zool.*, 30 A (23): 1-24.
- ANONYMOUS (1995): *Diagnóstico para la implementación de la Reserva de la Biosfera - Territorio Indígena, Pilón-Lajas*. Veterinarios sin Fronteras, La Paz. 184 pp.
- APARICIO, J. (1992): Herpetofauna, Pp. 113-119, in: Salm, H. & Marconi, M. (eds.): *Reserva Nacional Amazónica Manuripi-Heath. Programa de reestructuración (Fase II)*. Lidema, La Paz.
- APARICIO, J. (1993): Herpetofauna de Huara-co, un ecosistema andino en el altiplano central de Bolivia. *Ecol. Bol. Doc. ser. Zool.*, 4: 1-38.
- APARICIO, J. (1996): Herpetofauna. Pp 169-172, in: Plan de Manejo del Área Natural de Manejo Integral "Parque Nacional Sajama". VAIPO-DGB-GTZ, La Paz (unpublished).
- APARICIO, J. (1996): Herpetofauna de la carretera San Borja-Trinidad, Departamento del Beni (Bolivia). *Rev. Bol. de Ecol.* 1 (1): 53-60.
- APARICIO, J. (1999): Herpetofauna. Pp. 66-70, in: Estudio de Biodiversidad de la Reserva Nacional de Flora y Fauna Tariquía (Tarija). IE-FUNDECO, La Paz (unpublished).
- APARICIO, J. (1999): Herpetofauna de la Reserva Nacional de Fauna Altoandina Ulla-Ulla, Departamento de La Paz (Bolivia). *Rev. Ecol. Cons. Amb. Fund. Simón Patiño* (in press).
- BARBOUR, T. & NOBLE, G. K. (1920): Some amphibians from northwestern Peru with a revision of the genera *Phyllobates* and *Telmatobius*. *Bull. Mus. Comp. Zool.*, 63 (8): 394-433.
- BARRIO, A. (1965): Las subespecies de *Hyla pulchella* Duméril y Bibron (Anura, Hyliidae). *Physis*, 69: 115-128.
- BOETTGER, O. (1891): Reptilien und Batrachier aus Bolivia. *Zool. Anz.*, 14: 343-347.
- BOKERMANN, W. C. A. (1964): Notes on tree frogs of the *Hyla marmorata* group with description of a new species (Amphibia, Hyliidae). *Senck. Biol.*, 45 (3/5): 243-254.
- BOSCH, J., DE LA RIVA, I. & MÁRQUEZ, R. (1996): The calling behaviour of *Lysapsus limellus* and *Pseudis paradoxa* (Amphibia: Anura: Pseudidae). *Folia Zool.*, 45 (1): 49-55.
- BOULENGER, G. A. (1882): *Catalogue of the Batrachia Salientia S. Ecaudata in the collection of the British Museum*. 2nd ed., British Museum, London. 503 pp.
- BOULENGER, G. A. (1887): Description of

- new or little-known South American frogs of the genera *Paludicola* and *Hyla*. *Ann. Mag. Nat. Hist.*, (ser. 5) 20: 295-300.
- BOULENGER, G. A. (1891): Notes on American batrachians. *Ann. Mag. Nat. Hist.*, (ser. 8) 48: 453-457.
- BOULENGER, G. A. (1898): A list of the reptiles and batrachians collected by the late Prof. L. Balzan in Bolivia. *Ann. Mus. Civ. Stor. Nat. Genova*, (ser. 2) 19: 128-133.
- BOULENGER, G. A. (1902): Description of new batrachians and reptiles from the Andes of Peru and Bolivia. *Ann. Mag. Nat. Hist.*, (ser. 7) 10: 394-402.
- BOULENGER, G. A. (1903a): Exhibition of a specimen of *Ceratohyla bubalus* Espada, carrying eggs on its back. *Proc. Zool. Soc. London* 2 (1): 115-116.
- BOULENGER, G. A. (1903): Descriptions of new batrachians in the British Museum. *Ann. Mag. Nat. Hist.*, (ser. 7) 12: 552-557.
- BOULENGER, G. A. (1905): Description of new tailless batrachians in the collection of the British Museum. *Ann. Mag. Nat. Hist.*, (ser. 7) 16: 180-184.
- BRAME, A. H. & WAKE, D. B. (1963): The salamanders of South America. *Contr. Sci.*, 69: 5-71.
- CADLE, J. E. & PATTON, J. L. (1988): Distribution patterns of some amphibians, reptiles and mammals of the eastern Andean slope of southern Peru, pp. 225-244, in: Heyer, W. R. & Vanzolini, P. E. (eds.): *Proceedings of a workshop on Neotropical distribution patterns*. Academia Brasileira de Ciencias, Rio de Janeiro.
- CANNATELLA, D. C. (1980): Two new species of *Centrolenella* from Bolivia (Anura: Centrolenidae). *Proc. Biol. Soc. Wash.*, 93 (3): 714-724.
- CANNATELLA, D. C. (1983): Synonymy and distribution of *Phyllomedusa boliviiana* Boulenger (Anura: Hylidae). *Proc. Biol. Soc. Wash.*, 96 (1): 59-66.
- CANNATELLA, D. C. & DUELLMAN, W. E. (1982): Two new species of *Centrolenella*, with a brief review of the genus in Peru and Bolivia. *Herpetologica*, 38 (3): 380-388.
- CANNATELLA, D. C. & DUELLMAN, W. E. (1984): Leptodactylid frogs of the *Physalaemus pustulosus* group. *Copeia*, 1984 (4): 902-921.
- CANNATELLA, D. C., HILLIS, D. M., CHIPPIN-DALE, P. T., WEIGHT, L., RAND, A. S. & RYAN, M. J. (1998): Phylogeny of frogs of the *Physalaemus Pustulosus* [sic] species group, with an examination of data incongruence. *Syst. Biol.*, 47 (2): 311-335.
- CARAMASCHI, U. & DA CRUZ, C. A. G. (1997): Redescription of *Chiasmocleis albopunctata* (Boettger) and description of a new species of *Chiasmocleis* (Anura: Microhylidae). *Herpetologica*, 53 (2): 259-268.
- CARRIZO, G. R. (1992): Cuatro especies nuevas de anuros (Bufonidae: *Bufo* e Hyliidae: *Hyla*) del norte de la Argentina. *Cuad. Herp.*, 7 (3): 14-23.
- CEI, J. M. (1962): Mapa preliminar de la distribución continental de las "sibling species" del grupo *ocellatus* (género *Leptodactylus*). *Rev. Soc. Argent. Biol.* 38: 258-265.
- CEI, J. M. (1968): A new frog of the genus *Syrrhophus* from the Bolivian plateau. *J. Herpetol.*, 2 (3/4): 137-141.
- CEI, J. M. (1968): Remarks on the geographical distribution and phyletic trends of South American toads. *Pearce-Sellards Ser.*, 13: 1-21.
- CEI, J. M. (1972): *Bufo* of South America, pp. 82-92, in: Blair, W. F. (ed.): *Evolution in the genus Bufo*. Univ. Texas Press, Austin.
- CEI, J. M. (1980): Amphibians of Argentina. *Monogr. Monitore Zool. Ital.*, (2): 1-609.
- CEI, J. M. (1986): Speciation and adaptive radiation in Andean *Telmatobius* frogs, pp. 374-386, in: F. Vuilleumier and M. Monasterio (eds.), *High Altitude Tropical Biogeography*, Oxford Univ. Press.
- CEI, J. M. (1987): Additional notes to "Amphibians of Argentina": an update 1980-86. *Monitore Zool. Ital. (N.S.)*, 21: 209-272.

- CEI, J. M. (1990): On a Paraguayan sample of a long time confused species: *Physalaemus fuscomaculatus* (Steindachner, 1864) (Anura, Leptodactylidae). *Boll. Mus. reg. Sci. nat. Torino*, 8 (1): 215-231.
- CEI, J. M. & ERSPAMER, V. (1966): Biochemical taxonomy of South American amphibians by means of skin amines and polypeptides. *Copeia*, 1966 (1): 74-78.
- CEI, J. M., ERSPAMER, V. & ROSEGHINI, M. (1968): Taxonomic and evolutionary significance of biogenic amines and polypeptides in amphibian skin. II. Toads of the genera *Bufo* and *Melanophryniscus*. *Syst. Zool.*, 17 (3): 232-245.
- CÉSPEDEZ, J. A. (1997): Geographic distribution: *Pleurodema borelli*. *Herp. Rev.*, 28 (4): 208.
- CHRISTMANN, S. P. (1995): *Pleurodema cinereum* [sic], eine schaumnestproduzierende Krote aus Bolivien. *Elaphe N.F.*, 3 (4): 2-7.
- COCHRAN, D. (1955): Frogs of southeastern Brazil. *Bull. U.S. Nat. Mus.*, 206: 1-423.
- COCHRAN, D. & GOIN, C. J. (1970): Frogs of Colombia. *Bull. U.S. Nat. Mus.*, 288: 1-655.
- DE GRYS, P. (1938): *Hyla Roeschmanni* spec. nov. *Zool. Anz.*, 124: 315-316.
- DE LA RIVA, I. (1990a): Lista preliminar comentada de los anfibios de Bolivia con datos sobre su distribución. *Boll. Mus. reg. Sci. nat. Torino*, 8 (1): 261-319.
- DE LA RIVA, I. (1990b): Una especie nueva de *Oolygon* (Anura: Hylidae) procedente de Bolivia. *Rev. Esp. Herp.*, 4: 81-86.
- DE LA RIVA, I. (1992a): Comentarios sobre el género *Gastrotheca* (Anura: Hylidae) en Bolivia y descripción de una nueva especie. *Rev. Esp. Herp.*, 6: 15-22.
- DE LA RIVA, I. (1992b): A new species of *Phrynopus* from Bolivia (Anura: Leptodactylidae). *Herpetologica*, 48 (1): 111-114.
- DE LA RIVA, I. (1993a): *Ecología de una comunidad neotropical de anfibios durante la estación lluviosa*. Ph.D. thesis Univ. Complutense, Madrid. 365pp. (unpublished).
- DE LA RIVA, I. (1993b): A new species of *Sci-*
nax (Anura: Hylidae) from Argentina and Bolivia. *J. Herpetol.*, 27 (1): 41-46.
- DE LA RIVA, I. (1993c): Sinopsis del género *Eleutherodactylus* (Amphibia, Anura, Leptodactylidae) en Bolivia y adición de tres especies nuevas para el país. *Rev. Esp. Herp.*, 7: 97-105.
- DE LA RIVA, I. (1993d): The taxonomic status of *Hyla carinata* Andersson, 1938. *Herpetol. J.*, 3: 113-114.
- DE LA RIVA, I. (1994a): An undescribed defensive mechanism in the Neotropical hylid frog *Phrynohyas coriacea*. *Amphibia-Reptilia*, 15: 226-227.
- DE LA RIVA, I. (1994b): Geographic distribution: *Eleutherodactylus peruvianus*. *Herp. Rev.*, 25 (4): 159.
- DE LA RIVA, I. (1994c): A new aquatic frog of the genus *Telmatobius* (Anura: Leptodactylidae) from Bolivian cloud forests. *Herpetologica*, 50 (1): 38-45.
- DE LA RIVA, I. (1995a): Geographic distribution: *Melanophryniscus rubriventris*. *Herp. Rev.*, 26 (3): 152-153.
- DE LA RIVA, I. (1995b): A new reproductive mode for the genus *Adenomera* (Amphibia: Anura: Leptodactylidae): taxonomic implications for certain Bolivian and Paraguayan populations. *Stud. Neotrop. Fauna Environ.*, 30 (1): 15-29.
- DE LA RIVA, I. (1995c): Description of a new species of *Telmatobius* from Bolivia (Anura: Leptodactylidae). *Graellsia*, 50: 161-164.
- DE LA RIVA, I. (1995d): Geographic distribution: *Chiasmocleis ventrimaculata*. *Herp. Rev.*, 26 (2): 104.
- DE LA RIVA, I. (1996): The specific name of *Adenomera* (Anura: Leptodactylidae) in the Paraguay River Basin. *J. Herpetol.*, 30 (4): 556-558.
- DE LA RIVA, I. (1997): Redescription, variation, biology, and distribution of *Eleutherodactylus platydactylus* (Boulenger, 1903) (Amphibia: Anura: Leptodactylidae), the taxonomic status of *E. bockermannii* Donoso-Barros, 1970, and comments on poly-

- morphism in anurans. *Rev. Esp. Herp.*, 11: 71-90.
- DE LA RIVA, I. (1999a): First record of *Dendrophryniscus minutus* (Melin, 1941) in Bolivia (Anura: Bufonidae). *Herpetozoa*, 12 (1/2): 91-92.
- DE LA RIVA, I. (1999b): Geographic distribution: *Hyla schubarti*. *Herp. Rev.*, 30 (2): 107.
- DE LA RIVA, I. (1999c): Geographic distribution: *Scarthyla ostinodactyla*. *Herp. Rev.*, 30 (2): 108.
- DE LA RIVA, I. (1999d): Notes on the Paradox Frog, *Pseudis paradoxa*, in Bolivia. *Brit. Herp. Soc. Bull.*, 68: 14-19.
- DE LA RIVA, I. (1999): A new *Phyllomedusa* from southwestern Amazonia (Amphibia: Anura: Hylidae). *Rev. Esp. Herp.*, 13: 123-131.
- DE LA RIVA, I., BOSCH, J. & MÁRQUEZ, R. (1996a): Advertisement calls of two Bolivian toads (Anura: Bufonidae: *Bufo*). *Herpetol. J.*, 6: 59-61.
- DE LA RIVA, I., CASTROVIEJO, J. & CABOT, J. (1992): *Pseustes sulphureus* (Wagler, 1824) (Serpentes: Colubridae) nueva especie para Bolivia y datos sobre la herpetofauna boliviana. *Acta zool. Lilloana*, 41: 215-218.
- DE LA RIVA, I. & DUELLMAN, W. E. (1997): The identity and distribution of *Hyla rossaleni* Goin. *Amphibia-Reptilia*, 18: 433-436.
- DE LA RIVA, I. & GONZÁLES, L. (1998). Sobre la presencia de *Pleurodema guayapae* Barrio, 1964 (Anura, Leptodactylidae) en Bolivia. *Alytes*, 16 (1/2): 68-76.
- DE LA RIVA, I. & KÖHLER, J. (1998): A new minute leptodactylid frog, genus *Phyllonastes*, from humid montane forests of Bolivia. *J. Herpetol.*, 32 (3): 325-329.
- DE LA RIVA, I. & LYNCH, J. D. (1997): New species of *Eleutherodactylus* from Bolivia (Amphibia: Leptodactylidae). *Copeia*, 1997 (1): 151-157.
- DE LA RIVA, I. & MALDONADO, M. (1999): First record of *Leptodactylus ocellatus* (Linnaeus, 1758) in Bolivia (Anura, Leptodactylidae) and comments on related forms. *Graellsia*, 55: 193-197.
- DE LA RIVA, I., MÁRQUEZ, R. & BOSCH, J. (1994): Advertisement calls of Bolivian species of *Scinax* (Amphibia, Anura, Hylidae). *Bijdr. Dierk.*, 64 (2): 75-85.
- DE LA RIVA, I., MÁRQUEZ, R. & BOSCH, J. (1995): Advertisement calls of eight Bolivian hylids (Amphibia: Anura). *J. Herpetol.*, 29 (1): 113-118.
- DE LA RIVA, I., MÁRQUEZ, R. & BOSCH, J. (1996b): The advertisement calls of three South American poison frogs (Amphibia: Anura: Dendrobatiidae), with comments on their taxonomy and distribution. *J. Nat. Hist.*, 30: 1413-1420.
- DE LA RIVA, I., MÁRQUEZ, R. & BOSCH, J. (1996c): Advertisement calls of four micro-hylid frogs from Bolivia (Amphibia, Anura). *Am. Midl. Nat.*, 136 (2): 418-422.
- DE LA RIVA, I., MÁRQUEZ, R. & BOSCH, J. (1997): Description of the advertisement calls of some South American Hylidae (Amphibia, Anura): taxonomic and methodological consequences. *Bonn. zool. Beitr.*, 47 (1/2): 175-185.
- DE MACEDO, H. (1960): Vergleichende Untersuchungen an Arten der Gattung *Telmatobius* (Amphibia, Anura). *Zeitsch. Wissenschaft. Zool.*, 163 (3/4): 355-396.
- DE SÁ, R. O. (1995): *Hyla albopunctata*. *Cat. Amer. Amphib. Rept.*: 602.1-602.5.
- DONOSO-BARROS, R. (1969): Una nueva especie de Atelopodidae de Bolivia (Anura). *Physis*, 28: 327-330.
- DONOSO-BARROS, R. (1969): Un nuevo anuro de Bolivia, *Eupemphix freibergi* nov. sp. *Bol. Soc. Biol. Concepción*, 41: 183-187.
- DONOSO-BARROS, R. (1970): Nueva rana del genero *Eleutherodactylus* de la Amazonia Boliviana. *Bol. Soc. Biol. Concepción*, 42: 17-21.
- D'ORBIGNY, A. (1847): *Voyage à la Amérique Meridionale*. Vol. 5. P. Bertrand, Paris.
- DUELLMAN, W. E. (1956): The frogs of the hylid genus *Phrynohyas* Fitzinger, 1843. *Misc. Publ. Mus. Zool. Univ. Michigan*, 96: 1-59.

- DUELLMAN, W. E. (1970): The hylid frogs of Middle America. *Monogr. Mus. Nat. Hist. Univ. Kansas*, 1: 1-753.
- DUELLMAN, W. E. (1971): A taxonomic review of South American hylid frogs, genus *Phrynohyas*. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 4: 1-21.
- DUELLMAN, W. E. (1972): South American frogs of the *Hyla rostrata* group (Amphibia, Anura, Hylidae). *Zool. Meded. Leiden*, 47 (14): 179-195.
- DUELLMAN, W. E. (1972): The systematic status and life history of *Hyla rhodopepla* Günther. *Herpetologica*, 28 (4): 369-375.
- DUELLMAN, W. E. (1973): Frogs of the *Hyla geographica* group. *Copeia*, 1973 (3): 515-533.
- DUELLMAN, W. E. (1974): A reassessment of the taxonomic status of some Neotropical hylid frogs. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 27: 1-27.
- DUELLMAN, W. E. (1974): Taxonomic notes on *Phyllomedusa* (Anura, Hylidae) from the upper Amazon Basin. *Herpetologica*, 30 (2): 105-112.
- DUELLMAN, W. E. (1977): Liste der rezenten Amphibien und Reptilien: Hylidae, Centrolenidae, Pseudidae. *Das Tierreich*, 95: 1-225.
- DUELLMAN, W. E. (1978): New species of leptodactylid frogs of the genus *Eleutherodactylus* from the Cosñipata Valley, Peru. *Proc. Biol. Soc. Wash.*, 91 (2): 418-430.
- DUELLMAN, W. E. (1979): The herpetofauna of the Andes: patterns of distribution, origin, differentiation and present communities, pp. 371-459, in: Duellman, W. E. (ed.): *The South American herpetofauna: its origin, evolution, and dispersal*. *Mus. Nat. Hist. Monogr. Univ. Kansas* 7.
- DUELLMAN, W. E. (1982): Quaternary climatic-ecological fluctuation in the lowland tropics: frogs and forests, pp. 641-657, in: Prance, G. T. (ed.): *Biological diversification in the tropics*. Columbia University Press, New York.
- DUELLMAN, W. E. (1993): Amphibian species of the world: additions and corrections. *Univ. Kansas Spec. Publ.*, 21: 1-372.
- DUELLMAN, W. E. (1999): Distribution patterns of amphibians in South America., pp. 255-328, in: Duellman, W. E. (ed): *Patterns of distribution of amphibians. A global perspective*. The John Hopkins University Press, Baltimore & London.
- DUELLMAN, W. E. & DE LA RIVA, I. (1999): Rediscovery and taxonomic status of *Hyla splendens* Schmidt, 1857 (Anura, Hylidae). *Copeia*, 1999 (1): 197-199.
- DUELLMAN, W. E., DE LA RIVA, I. & WILD, E. R. (1997): Frogs of the *Hyla armata* and *Hyla pulchella* groups in the Andes of South America, with definitions and analyses of phylogenetic relationships of Andean groups of *Hyla*. *Sci. Pap. Nat. Hist. Mus. Univ. Kansas*, 3: 1-41.
- DUELLMAN, W. E. & FRITTS, T. H. (1972): A taxonomic review of the southern Andean marsupial frogs (Hylidae: *Gastrotheca*). *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 9: 1-37.
- DUELLMAN, W. E. & SCHULTE, R. (1992): Description of a new species of *Bufo* from northern Peru with comments on phenetic groups of South American toads (Anura: Bufonidae). *Copeia*, 1992 (1): 162-172.
- DUELLMAN, W. E. & THOMAS, R. (1996): Anuran amphibians from a seasonally dry forest in southeastern Peru and comparison of the anurans among sites in the upper Amazon Basin. *Occ. Pap. Nat. Hist. Mus. Univ. Kansas*, 180: 1-34.
- DUELLMAN, W. E. & TRUEB, L. (1986): *Biology of amphibians*. McGraw-Hill, New York. 670 pp.
- DUELLMAN, W. E. & VELOSO, A. (1977): Phylogeny of *Pleurodema* (Anura: Leptodactylidae): a biogeographic model. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 64: 1-46.
- DUELLMAN, W. E. & WIENS, J. J. (1993): Hylid frogs of the genus *Scinax* Wagler, 1830, in Amazonian Ecuador and Peru. *Occ.*

- Pap. Mus. Nat. Hist. Univ. Kansas*, 153: 1-57.
- DUMÉRIL, A. M. C. & BIBRON, G. (1841): *Erpétologie générale ou histoire naturelle compléte des reptiles*. Vol. 8, Paris. 792 pp.
- DUNN, E. R. (1942): The American caecilians. *Bull. Mus. Comp. Zool.*, 91 (6): 439-550.
- DUNN, E. R. (1949): Notes on South American frogs of the family Microhylidae. *Amer. Mus. Novit.*, 1419: 1-21.
- EDWARDS, S. R. (1974): *A phenetic analysis of the genus Colostethus (Anura, Dendrobatidae)*. Ph.D. Thesis, Univ. Kansas. 420 pp. (unpublished).
- EISENTRAUT, M. (1932): Biologische Beobachtungen im bolivianischen Chaco. V. Ein neuer Fall von Brutfürsorge in der Froschgattung *Leptodactylus*. *Zeitschr. Morph. Ökol. Tiere*, 26 (1/2): 317-326.
- EMMONS, L. H. (1991): Observations on the herpetofauna, p. 74, in: Parker, T. A. & Bailey, B. (eds.): *A biological assessment of the Alto Madidi region and adjacent areas of northwest Bolivia*. RAP Working Papers 1, Conservation International, Washington.
- EMMONS, L. H. (1993): Reptile and amphibian species recorded in the Curuyuqui Chaco forest, p. 104, in: Parker, T. A., III, Gentry, A. H., Foster, R. B., Emmons, L. H., & Remsen, J. V. Jr. (eds.): *The lowland dry forests of Santa Cruz, Bolivia: a global conservation priority*. RAP Working Papers 4, Conservation International, Washington.
- ERGUETA, P. (1991): Los Anfibios, pp. 453-468, in: Forno, E. & Baudo, M. (eds.): *Historia natural de un valle de los Andes: La Paz*. Instituto de Ecología, UMSA, La Paz.
- ERGUETA, P. (1992): Panorama de la herpetología en Bolivia. *Acta zool. Lilloana*, 41: 5-12.
- ERGUETA, P. (1993): Aspectos de la biología y ecología de *Phrynobatrachus laplacai* (Anura: Leptodactylidae) en un bosque nublado de altura de Yungas (La Paz, Bolivia). *Ecología en Bolivia*, 21: 19-29.
- ERGUETA, P. & GÓMEZ, H. (eds.). (1997): *Directorio de áreas protegidas de Bolivia*. Centro de Datos para la Conservación, La Paz. 186 pp.
- ERGUETA, P. & HARVEY, M. B. (1996): Anfibios, pp. 67-72, in: Ergueta, P. & Morales, C. (eds.): *Libro rojo de los vertebrados de Bolivia*. Centro de Datos para la Conservación, La Paz.
- ERGUETA, P. & MORALES, C. (eds.) (1996): *Libro rojo de los vertebrados de Bolivia*. Centro de Datos para la Conservación, La Paz. 347 pp.
- ETZEL, V. V. (1988): Die Schwarzfleckenkröte aus dem nördlichen Argentinien. *Melanophryniscus rubriventris*. *D. Aquar. Z. (DATZ)*, 41 (9): 348-350.
- FROST, D. R. (ed.) (1985): *Amphibian species of the world. A taxonomic and geographical reference*. Allen Press & ASC, Lawrence, Kansas, USA. 732 pp.
- FUGLER, C. M. (1983): Lista preliminar de los anfibios y de los reptiles de Tumi Chucua, Provincia Vaca Díez, Departamento de Beni. *Mus. Nac. Hist. Nat. (Bolivia) Comunicación*, 2: 4-11.
- FUGLER, C. M. (1984): Tercera contribución a la fauna herpetológica del Oriente Boliviano. *Ecología en Bolivia*, 5: 63-72.
- FUGLER, C. M. (1985): Adiciones y correcciones a la lista preliminar de la herpetofauna de Tumi Chucua, Provincia Vaca Díez, Departamento de Beni. *Cuad. Acad. Cienc. Bolivia (Zool.)*, 3: 17-18.
- FUGLER, C. M. (1986): La estructura de una comunidad herpetológica en las selvas benianas en la estación de sequía. *Ecología en Bolivia*, 8: 1-20.
- FUGLER, C. M. (1988): Consideraciones sobre la herpetofauna de las selvas tropicales bolivianas. *Mus. Nac. Hist. Nat. (Bolivia) Comunicación*, 6: 3-12.
- FUNKHOUSER, A. (1957): A review of the Neotropical tree frogs of the genus *Phyllomedusa*. *Occ. Pap. Nat. Hist. Mus. Stanford Univ.*, 5: 1-90.

- GAIGE, H. T. (1929): Three new tree-frogs from Panama and Bolivia. *Occ. Pap. Mus. Zool. Univ. Michigan*, 207: 1-6.
- GALLARDO, J. M. (1961a): Three new toads from South America: *Bufo manicorensis*, *Bufo spinulosus altiperuvianus* and *Bufo quechua*. *Breviora*, 141: 1-11.
- GALLARDO, J. M. (1961b): On the species of Pseudidae (Amphibia, Anura). *Bull. Mus. Comp. Zool.*, 125 (4): 111-134.
- GALLARDO, J. M. (1964a): Consideraciones sobre *Leptodactylus ocellatus* (L.) (Amphibia, Anura) y especies aliadas. *Physis*, B 24 (68): 373-384.
- GALLARDO, J. M. (1964b): "Leptodactylus prognathus" Boul. y "L. mystacinus" (Burm.) con sus respectivas especies aliadas. (Amphibia, Leptodactylidae del grupo «cavicola»). *Rev. Mus. Arg. Cienc. Nat. "Bernardino Rivadavia"*, 9 (5): 91-121.
- GALLARDO, J. M. (1964c): Una nueva forma de Pseudidae (Amphibia, Anura), y algunas consideraciones sobre las especies argentinas de esta familia. *Acta zool. Lilloana*, 20: 193-210.
- GALLARDO, J. M. (1965): The species *Bufo granulosus* Spix (Salentia: Bufonidae) and its geographic variation. *Bull. Mus. Comp. Zool.*, 134 (4): 107-138.
- GALLARDO, J. M. (1967): Un nuevo nombre para *Bufo granulosus minor* Gallardo. *Neotropica*, 13 (41): 56.
- GALLARDO, J. M. (1979): Composición, distribución y origen de la herpetofauna chaqueña, pp. 299-307, in: Duellman, W. E. (ed.): *The South American herpetofauna: its origin, evolution, and dispersal*. Univ. Kansas, Mus. Nat. Hist. Monogr. 7.
- GALLARDO, J. M. (1987): *Anfibios argentinos: guía para su identificación*. Librería Agropecuaria S.A., Buenos Aires. 98 pp.
- GANS, C. (1960): Notes on a herpetological collecting trip through the southeastern lowlands of Bolivia. *Ann. Carnegie Mus.*, 35: 283-314.
- GARMAN, S. W. (1875): Exploration of Lake Titicaca by Alexander Agassiz and S. W. Garman. Batrachians, reptiles. *Bull. Mus. Comp. Zool.*, 3: 276-278.
- GAVETTI, E. & ANDREONE, F. (1993): *Revised catalogue of the herpetological collection in Turin University. I. Amphibia*. Museo Regionale di Scienze Naturali, Torino. 185 pp.
- GONZÁLES, L. & FERNÁNDEZ, I. (1995): La herpetofauna de la Laguna Volcán, Provincia Florida, Departamento Santa Cruz-Bolivia: resultados preliminares. UAGRM-MHNNKM-BOLFOR (unpublished).
- GONZÁLES, L. (1998): La herpetofauna del Izozog. *Ecología en Bolivia*, 31: 45-51.
- GONZÁLES, L., LÖTTERS, S. & REICHLE, S. (1999): On the dendrobatid frogs from Bolivia, rediscovery of *Epipedobates boliviensis* (Boulenger, 1902), first record of *Colostethus brunneus* (Cope, 1887) and comments on other species (Anura: Dendrobatidae). *Herpetozoa* 12 (3/4): 179-186.
- GORHAM, S. W. (1966): Liste der rezenten Amphibien und Reptilien. *Das Tierreich*, 85: 1-222.
- GORIZULA, S. & SEÑARIS, J. C. (1998): Contribution to the herpetofauna of the Venezuelan Guayana. I. A data base. *Scientia Guiana*, 8: 1-268 + 32 plates.
- HADDAD, C. F. B. & MARTINS, M. (1994): Four species of Brazilian poison frogs related to *Epipedobates pictus* (Dendrobatidae): taxonomy and natural history observations. *Herpetologica*, 50 (3): 282-295.
- HARDING, K. A. (1983): *Catalogue of New World amphibians*. Pergamon Press. 406 pp.
- HARVEY, M. B. (1995): A preliminary list of the reptiles and amphibians of the El Refugio biological reserve, pp. 21-26, in: Forsyth, A. (ed.): *A report on aspects of biodiversity and conservation potential in El Refugio*. Conservation International, Washington.
- HARVEY, M. B. (1996): A new species of glass frog (Anura: Centrolenidae: *Cochranella*) from Bolivia, and the taxonomic status of *Cochranella flavidigitata*. *Herpetologica*, 52 (3): 427-435.

- HARVEY, M. B. (1997): Reptiles and amphibians from the vicinity of El Palmar in the Andes of Chuquisaca, Bolivia, pp. 33-36, 83-84, *in*: Schulenberg, T. S. & Awbrey, K. (eds.): *A rapid assessment of the humid forests of south central Chuquisaca, Bolivia*. RAP Working Papers 8, Conservation International, Washington.
- HARVEY, M. B. (1998): Reptiles and amphibians of Parque Nacional Noel Kempff Mercado, pp. 144-153, *in*: Killeen, T. J. & Schulenberg, T. S. (eds.): *A biological assessment of Parque Nacional Noel Kempff Mercado, Bolivia*. RAP Working Papers 10, Conservation International, Washington.
- HARVEY, M. B., APARICIO, J., CORTEZ, C., GONZÁLES, L., GUERRA, J. F., MONTAÑO, M. S. & PÉREZ, M. E. (1998): Reptile and amphibian species of Parque Nacional Noel Kempff Mercado, pp. 348-355, *in*: Killeen, T. J. & Schulenberg, T. S. (eds.): *A biological assessment of Parque Nacional Noel Kempff Mercado, Bolivia*. RAP Working Papers 10, Conservation International, Washington.
- HARVEY, M. B. & ERGUETA, P. (1998): A new species of *Phrynoporus* (Anura: Leptodactylidae) from elfin forest in the Bolivian Andes. *J. Herpetol.*, 32 (4): 536-540.
- HARVEY, M. B. & KECK, B. (1995): A new species of *Ischnocnema* (Anura: Leptodactylidae) from high elevations in the Andes of Central Bolivia. *Herpetologica*, 51 (1): 56-66.
- HARVEY, M. B. & SMITH, E. N. (1993): A new species of aquatic *Bufo* (Anura: Bufonidae) from cloud forests in the Serranía de Siberia, Bolivia. *Proc. Biol. Soc. Wash.*, 106 (3): 442-449.
- HARVEY, M. B. & SMITH, E. N. (1994): A new species of *Bufo* (Anura: Bufonidae) from cloud forests in Bolivia. *Herpetologica*, 50 (1): 32-38.
- HASS, C. A., DUNSKI, J. F., MAXSON, L. R. & HOOGMOED, M. S. (1995): Divergent lineages within the *Bufo margaritifera* Complex (Amphibia: Anura; Bufonidae) revealed by albumin immunology. *Biotropica*, 27 (2): 238-249.
- HÄUPL, M. & TIEDEMANN, F. (1978): Typenkatalog der herpetologischen Sammlung. *Katal. wiss. Samml. Naturhist. Mus. Wien Vertebrata*, 2: 1-35.
- HENLE, K. (1992): Zur Amphibienfauna Perus nebst Beschreibung eines neuen *Eleutherodactylus* (Leptodactylidae). *Bonn. zool. Beitr.*, 43 (1): 79-129.
- HEYER, W. R. (1970): Studies on the frogs of the genus *Leptodactylus* (Amphibia, Leptodactylidae). VI. Biosystematics of the *melanonotus* group. *Cont. Sci. Nat. Hist. Los Angeles Co.*, 191: 1-48.
- HEYER, W. R. (1973): Systematics of the *marmoratus* group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Cont. Sci. Nat. Hist. Los Angeles Co.*, 251: 1-50.
- HEYER, W. R. (1977): Taxonomic notes on frogs from the Madeira and Purus rivers, Brasil. *Pap. Avul. Zool. São Paulo*, 31 (8): 141-162.
- HEYER, W. R. (1978): Systematics of the *fuscus* species group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Bull. Los Angeles Co. Mus. Nat. Hist.*, 29: 1-85.
- HEYER, W. R. (1979): Systematics of the *pentadactylus* species group of the frog genus *Leptodactylus* (Amphibia, Leptodactylidae). *Smiths. Contrib. Zool.*, 301: 1-43.
- HEYER, W. R. (1994): Variation within the *Leptodactylus podicipinus-wagneri* complex of frogs (Amphibia: Leptodactylidae). *Smiths. Contrib. Zool.*, 546: 1-124.
- HEYER, W. R., GARCÍA-LÓPEZ, J. M. & CARDOSO, A. J. (1996): Advertisement call variation in the *Leptodactylus mystaceus* species complex (Amphibia: Leptodactylidae) with a description of a new sibling species. *Amphibia-Reptilia*, 17: 7-31.
- HEYER, W. R. & MUÑOZ, A. M. (1999): Validation of *Eleutherodactylus crepitans* Bokermann, 1965, notes on the types and type

- locality of *Telatrema heterodactylum* Miranda-Ribeiro, 1937, and description of a new species of *Eleutherodactylus* from Mato Grosso, Brazil (Amphibia: Anura: Leptodactylidae). *Proc. Biol. Soc. Wash.*, 112 (1): 1-18.
- HILLIS, D. M. & DE SÁ, R. O. (1988): Phylogeny and taxonomy of the *Rana palmipes* group (Salientia, Ranidae). *Herpetol. Monogr.*, 2: 1-26.
- HOOGMOED, M. S. (1979): The herpetofauna of the Guianan region, pp. 241-279, in: Duellman, W.E. (ed.): *The South American herpetofauna: its origin, evolution, and dispersal*. Univ. Kansas, Mus. Nat. Hist. Monogr. 7.
- HOOGMOED, M. S. (1986): Biosystematic studies of the *Bufo "typhonius"* group. A preliminary progress report, pp. 147-150, in: Roček, Z. (ed.): *Studies in Herpetology*. Prague.
- HOOGMOED, M. S. (1989): South American bufonids (Amphibia: Anura: Bufonidae), an enigma for taxonomists, pp. 167-180, in: Fontanet, X. & Horta, N. (eds.): *Trebballs d'ictiologia i herpetologia*. Treb. Soc. Cat. Ictio. Herp. 2.
- HOOGMOED, M. S. (1990): Biosystematics of South American Bufonidae, with special reference to the *Bufo "typhonius"* group, pp. 113-123, in: Peters, G. & Hutterer, R. (eds.): *Vertebrates in the tropics*. Museum Alexander Koenig, Bonn.
- HOOGMOED, M. S. (1993): The herpetofauna of floating meadows, pp. 199-213, in: Ouboter, P. E. (ed.): *Freshwater ecosystems of Suriname*. Kluwer Academic Publishers.
- HOOGMOED, M. S. & VACA, N. (1990): Herpetology, pp. 1-23, in: Frey, R. (ed.): *Report Expedition Perseverancia, 1990* (unpublished).
- HUTTERER, R. (1997): Archaeozoological remains (Vertebrata, Gastropoda) from pre-hispanic sites at Pailón, Bolivia. *Beitr. Allg. Vergl. Arch.*, 17: 325-341.
- IBISCH, P. L. (1996): Neotropische Epiphyten- diversitat - das Beispiel Bolivien. *Arch. naturw. Diss. (M. Galunder Verlag)* 1: 1-356.
- IBISCH, P. & BÖHME, W. (1993): Zur Kenntnis der innerandinen Herpetofauna Boliviens (Provinz Arque, Departamento Cochabamba). *Herpetofauna*, 15 (84): 15-26.
- IBISCH, P., ROJAS, P., DE LA BARRA, N., FERNÁNDEZ, E., MERCADO, M., OVANDO, L. & VARGAS, G. (1996): Un "lugar de encuentro": Flora de la zona arqueológica "El Fuerte", Samaipata, (Prov. Florida, Dpto. Santa Cruz, Bolivia). *Ecología en Bolivia*, 28: 1-28.
- KLAPPENBACH, M. A. (1985): Notas herpetológicas, V. *Com. Zool. Mus. Hist. Nat. Montevideo*, 11 (150): 1-23.
- KLAPPENBACH, M. A. & LANGONE, J. A. (1992): Lista sistemática y sinonímica de los anfibios del Uruguay con comentarios y notas sobre su distribución. *An. Mus. Nac. Hist. Nat. Montevideo* 2^a ser. (8): 163-222.
- KÖHLER, J. (1995a): *Untersuchungen zur Taxonomie, Ökologie und Zoogeographie bolivianischer Froschlurche (Amphibia: Anura)*. Diplomarbeit Univ. Bonn: 1-123 (unpublished).
- KÖHLER, J. (1995b): Erstnachweis von *Hyla marmorata*, *Leptodactylus syphax* und *Phrynoporus peruvianus* für Bolivien. *Herpetofauna*, 17 (98): 6-8.
- KÖHLER, J. (1997a): Geographic distribution: *Chacophrys pierotti*. *Herp. Rev.*, 28 (3): 155-156.
- KÖHLER, J. (1997b): Geographic distribution: *Lepidobatrachus laevis*. *Herp. Rev.*, 28 (3): 156.
- KÖHLER, J. (2000a): *Amphibian diversity in Bolivia: a study with special reference to montane forest regions*. - Ph. D. Dissertation, Univ. Bonn: 1-281 (unpublished).
- KÖHLER, J. (2000b): New species of *Eleutherodactylus* (Anura: Leptodactylidae) from cloud forest of Bolivia. *Copeia*, 2000 (2): 516-520.
- KÖHLER, J. (2000c): A new species of *Phyllonastes Heyer* from the Chapare region of

- Bolivia, with notes on *Phyllonastes carasscoicola* (Amphibia: Anura: Leptodactylidae). *Spixiana* 23 (1): 47-53.
- KÖHLER, J. & BÖHME, W. (1996): Anuran amphibians from the region of Pre-Cambrian rock outcrops (inselbergs) in northeastern Bolivia, with a note on the gender of *Scinax* Wagler, 1830 (Hylidae). *Revue fr. Aquariol. Herpétop.*, 23 (3/4): 133-140.
- KÖHLER, J., DIRKSEN, L., IBISCH, P. L., RAUER, G., RUDOLPH, D. & BÖHME, W. (1995a): Zur Herpetofauna des Sehuencas-Bergregenwaldes im Carrasco-Nationalpark, Bolivien. *Herpetofauna*, 17 (96): 12-25.
- KÖHLER, J., IBISCH, P. L., DIRKSEN, L. & BÖHME, W. (1995b): Zur Herpetofauna der semihumiden Samaipata-Region, Bolivien. I. Amphibien. *Herpetofauna*, 17 (98): 13-24.
- KÖHLER, J. & JUNGFER, K.-H. (1995): Eine neue Art und ein Erstnachweis von Fröschen der Gattung *Eleutherodactylus* aus Bolivien. *Salamandra*, 31 (3): 149-156.
- KÖHLER, J. & LÖTTERS, S. (1999a): Annotated list of amphibian records from the Departamento Pando, Bolivia, with description of some advertisement calls. *Bonn. zool. Beitr.* 48 (3/4): 259-273.
- KÖHLER, J. & LÖTTERS, S. (1999b): Advertisement calls of two Bolivian *Leptodactylus* (Amphibia: Anura: Leptodactylidae). *Amphibia-Reptilia*, 20: 215-219.
- KÖHLER, J. & LÖTTERS, S. (1999c): New species of the *Eleutherodactylus unistriatus* group (Amphibia: Anura: Leptodactylidae) from montane rain forest of Bolivia. *Copeia*, 1999 (2): 422-427.
- KÖHLER, J. & LÖTTERS, S. (2000): Description of a small tree frog, genus *Hyla* (Anura: Hylidae), from humid Andean slopes of Bolivia. *Revue fr. Aquariol. Herpétop.* (in press).
- KÖHLER, J., LÖTTERS, S. & REICHLE, S. (1998b): Amphibian species diversity in Bolivia, pp. 329-335, in: Barthlott, W. & Winiger, M. (eds.): *Biodiversity - a challenge for development research and policy*. Springer, Berlin, Heidelberg, New York.
- KÖHLER, J., MORALES, V. R., LÖTTERS, S., REICHLE, S. & APARICIO, J. (1998a): A new green species of frog, genus *Eleutherodactylus*, from Bolivia and Perú (Amphibia, Anura, Leptodactylidae). *Stud. Neotrop. Fauna Environm.*, 33 (2/3): 93-99.
- KÖHLER, J. & REICHLE, S. (1998): Geographic distribution: *Cochranella pluvialis*. *Herp. Rev.*, 29 (3): 172.
- KÖHLER, J., REICHLE, S. & PETERS, G. (1997): Advertisement calls of three species of *Bufo* (Amphibia: Anura: Bufonidae) from lowland Bolivia. *Stuttg. Beitr. Naturk. (Ser. A)*, 562: 1-8.
- LA MARCA, E. (1992): Catálogo taxonómico, biogeográfico y bibliográfico de las ranas de Venezuela. *Cuadernos Geográficos* 9: 1-197.
- LANGONE, J. A. & BASSO, N. G. (1987): Distribución geográfica y sinonimia de *Hyla nana* Boulenger, 1889 y de *Hyla sanborni* Schmidt, 1944 (Anura: Hylidae) y observaciones sobre formas afines. *Comun. Zool. Mus. Hist. Nat. Montevideo*, 11 (164): 1-17.
- LAURENT, R. F. (1967): Redescubrimiento de *Pithecopus pailonus* (Shreve) en Argentina. *Acta zool. Lilloana*, 22: 231-247.
- LAURENT, R. F. (1969): Las supuestas razas de *Bufo arenarum* y el concepto de subespecie críptica. *Acta zool. Lilloana* 25 (6): 69-78.
- LAVILLA, E. O. (1988): Lower Telmatobiinae (Anura: Leptodactylidae): generic diagnoses based on larval characters. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 124: 1-19.
- LAVILLA, E. O. (1988): *Telmatobius* (Anura, Leptodactylidae): the name-bearing types of five Vellard's taxa. *Alytes*, 7 (1): 6-18.
- LAVILLA, E. O. (1992): Tipos portadores de nombre y localidades tipo de anfibios de Argentina. *Acta zool. Lilloana*, 42 (1): 61-100.
- LAVILLA, E. O., CRUZ, F. B. & SCROCCHI, G. J. (1995): Amphibiens et reptiles de la station biologique "Los Colorados" dans la

- province de Salta, Argentine. - *Revue fr. Aquariol. Herpétol.*, 22 (1/2): 51-58.
- LAVILLA, E. O. & DE LA RIVA, I. (1993): La larva de *Telmatobius boliviensis*. *Alytes*, 11 (2): 37-46.
- LAVILLA, E. O., DE SÁ, R. O. & DE LA RIVA, I. (1997): Description of the tadpole of *Atelopus tricolor*. *J. Herpetol.*, 31 (1): 121-124.
- LAVILLA, E. O. & ERGUETA, P. (1995a): Una nueva especie de *Telmatobius* (Anura: Leptodactylidae) del sudoeste de Bolivia. *Ecología en Bolivia*, 24: 91-101.
- LAVILLA, E. O. & ERGUETA, P. (1995b): Una nueva especie de *Telmatobius* (Anura, Leptodactylidae) de la ceja de montaña de La Paz (Bolivia). *Alytes*, 13 (2): 45-51.
- LAVILLA, E. O. & ERGUETA, P. (1999): A new Bolivian species of the genus *Telmatobius* (Anura: Leptodactylidae) with a humeral spine. *Amphibia-Reptilia*, 20: 55-64.
- LAVILLA, E. O., GONZÁLES, L. & FERNÁNDEZ, I. (1996): *Herpetología Amboroense. Informe sobre la herpetofauna del Parque Nacional Amboró y áreas aledañas*. Consorcio FAN-TNC, 46 pp. (unpublished).
- LAVILLA, E. O. & SCROCCHI, G. J. (1999): Anfibios y Reptiles de Tariquía, pp. 83-92, in: J.A. González, G.J. Scrocchi & E.O. Lavilla (eds.): *Relevamiento de la biodiversidad de la Reserva Nacional de Flora y Fauna Tariquía (Tarija, Bolivia)*. Ser. Conservación de la Naturaleza, 14. Fundación Miguel Lillo - PROBONA.
- LESCURE, J., MARTY, V., MARTY, C., STARACE, F., AUBER-THOMAY, M. & LETELIER, F. (1995): Contribution à l'étude des Amphibiens de Guyane française. X. Les *Phyllomedusa* (Anura, Hylidae). *Revue fr. Aquariol.* 22 (1/2): 35-50.
- LOBO, F. (1992): Distribución y lista de localidades de *Pseudopaludicola* (Anura: Leptodactylidae) en la República Argentina. *Cuad. Herp.*, 7 (5): 28-37.
- LOBO, F. (1993): Descripción de una nueva especie del género *Physalaemus* (Anura: Leptodactylidae) del noroeste argentino. *Rev. Esp. Herp.*, 7: 13-20.
- LOBO, F. (1993): Lista bibliográfica comentada de *Pseudopaludicola* (Anura: Leptodactylidae) hasta junio de 1991. *Acta zool. Lilloana*, 42 (2): 197-208.
- LOBO, F. (1994): Primera cita de *Pseudopaludicola boliviensis* Parker, 1927, para el Brasil y nuevos registros en Paraguay. *Cuad. Herp.*, 8 (2): 231-232.
- LÖTTERS, S. (1996): *The Neotropical toad genus Atelopus. Checklist, biology, distribution*. Vences & Glaw Verlag, Cologne. 143 pp.
- LÖTTERS, S. & DE LA RIVA, I. (1998): Redescription of *Atelopus tricolor* Boulenger from southeastern Peru and adjacent Bolivia, with comments on related forms. *J. Herpetol.*, 32 (4): 481-488.
- LÖTTERS, S., DE LA RIVA, I., REICHLE, S. & SOTO, G. (2000): First records of *Bufo guttatus* (Amphibia: Bufonidae) from Bolivia, with notes on *Bufo glaberrimus*. *Bonn. zool. Beitr.* 49 (1/2): (in press).
- LÖTTERS, S., GLAW, F., REICHLE, S., KÖHLER, J. & MEYER, E. (1999a): Notes on vocalizations in three species of *Atelopus* from Central and South America (Amphibia: Anura). *Herpetozoa*, 12 (1/2): 79-83.
- LÖTTERS, S. & KÖHLER, J. (2000a): A new toad of the *Bufo typhonius* complex from humid montane forests in Bolivia (Amphibia: Anura, Bufonidae). *Spixiana* (in press).
- LÖTTERS, S. & KÖHLER, J. (2000b): *Cochranella nola* (Amphibia: Anura: Centrolenidae): natural history notes, distribution, and advertisement call. *Herp. Nat. Hist.* (in press).
- LÖTTERS, S., KÖHLER, J. & REICHLE, S. (1999b): Description of the tadpole of the Andean tree frog *Hyla mariannae* (Amphibia: Anura: Hylidae). *Folia Zool.*, 48 (1): 49-53.
- LUTZ, B. (1949): Frogs in the Adolpho Lutz collection at the Instituto Oswaldo Cruz. IV. Species allied to the green *Hylas* from SE

- Brazil. Addendum to part II. *Mem. Inst. Oswaldo Cruz*, 47 (3/4): 319-335.
- LUTZ, B. (1973): *Brazilian species of Hyla*. Univ. Texas Press, Austin & London. 260 pp.
- LYNCH, J. D. (1975): A review of the Andean leptodactylid frog genus *Phrynobius*. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 35: 1-51.
- LYNCH, J. D. (1976): The species groups of the South American frogs of the genus *Eleutherodactylus* (Leptodactylidae). *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 61: 1-24.
- LYNCH, J. D. (1979): The amphibians of the lowland tropical rainforests, pp. 189-215, in: Duellman, W. E. (ed.): *The South American herpetofauna: its origin, evolution, and dispersal*. Univ. Kansas Mus. Nat. Hist. Monogr., 7.
- LYNCH, J. D. (1980): A taxonomic and distributional synopsis of the Amazonian frogs of the genus *Eleutherodactylus*. *Amer. Mus. Novit.*, 2696: 1-24.
- LYNCH, J. D. (1986): Origins of the high Andean herpetofauna, pp. 478-499, in: Vuilleumier, F. & Monasterio, M. (eds.): *High altitude tropical biogeography*. Oxford Univ. Press.
- LYNCH, J. D. (1989a): A review of the leptodactylid frogs of the genus *Pseudopaludicola* in northern South America. *Copeia*, 1989 (3): 577-588.
- LYNCH, J. D. (1989b): Intrageneric relationships of mainland *Eleutherodactylus* (Leptodactylidae). I. A review of the frogs assigned to the *Eleutherodactylus discoidalis* species group. *Milwaukee Publ. Mus. Contr. Biol. Geol.*, 79: 1-25.
- LYNCH, J. D. & DUELLMAN, W. E. (1997): Frogs of the genus *Eleutherodactylus* in western Ecuador. *Nat. Hist. Mus. Univ. Kansas Spec. Publ.*, 23: 1-236.
- LYNCH, J. D. & McDIARMID, R. W. (1987): Two new species of *Eleutherodactylus* (Amphibia: Anura: Leptodactylidae) from Bolivia. *Proc. Biol. Soc. Wash.*, 100 (2): 337-346.
- MÁRQUEZ, R., DE LA RIVA, I. & BOSCH, J. (1993): Advertisement calls of Bolivian species of *Hyla* (Amphibia, Anura, Hylidae). *Biotropica*, 25 (4): 426-443.
- MÁRQUEZ, R., DE LA RIVA, I. & BOSCH, J. (1995): Advertisement calls of Bolivian Leptodactylidae (Amphibia, Anura). *J. Zool., Lond.*, 237: 313-336.
- MÁRQUEZ, R., DE LA RIVA, I. & BOSCH, J. (1996): Advertisement calls of three glass frogs from the Andean forests (Amphibia: Anura: Centrolenidae). *Herpetol. J.*, 6: 97-99.
- MCLISTER, J. D., LOUGHEED, S. C. & BOGART, J. P. (1991): Electrophoretic and vocalization comparisons among three leptodactylid frogs (*Pleurodema* spp.) from northwestern Argentina. *Can. J. Zool.*, 69: 2397-2403.
- MERCADAL, I. T. (1986): *Ceratophrys joazeirensis* sp. n. (Ceratophryidae, Anura) del noroeste de Brasil. *Amphibia-Reptilia*, 7: 313-334.
- MERTENS, R. (1929): Herpetologische Mitteilungen. XII. Über einige Amphibien und Reptilien aus Süd-Bolivien. *Zool. Anz.*, 86: 57-62.
- MONTAÑO, M. E. (1995): La fauna herpetológica. Inventario de base en la Concesión Oquiriquia. Proyecto BOLFOR (unpublished).
- MORALES, V. R. (1994): Taxonomía sobre algunos *Colostethus* (Anura: Dendrobatidae) de Sudamérica, con descripción de dos especies nuevas. *Rev. Esp. Herp.*, 8: 95-103.
- MÜLLER, L. (1924): Ueber neue oder seltene Mittel- und Südamerikanische Amphibien und Reptilien. *Mitt. Zool. Mus. Berlin*, 11: 77-93.
- MÜLLER, L. & HELLMICH, W. (1936): *Wissenschaftliche Ergebnisse der deutschen Gran Chaco-Expedition. Amphibien und Reptilien. I. Teil: Amphibia, Chelonia, Loricata*. Verlag von Strecker & Schröder, Stuttgart, 120 pp.
- NAPOLI, M. F. & CARAMASCHI, U. (1999):

- Geographic variation of *Hyla rubricundula* and *Hyla anataliasiasi*, with the description of a new species (Anura, Hylidae). *Alytes*, 16 (3/4): 165-189.
- NIEDEN, F. (1923): Anura I. Subordo Aglossa und Phaneroglossa. Arcifera. *Das Tierreich*, 46: 1-584.
- PACHECO, L. (1990): Herpetofauna, pp. 165-168, in: Diagnóstico de los recursos naturales de la Reserva Nacional de Fauna Andina "Eduardo Avaroa", IE-MNHN-CEDI, La Paz (unpublished).
- PARKER, H. W. (1927): A revision of the frogs of the genera *Pseudopaludicola*, *Physalaemus* and *Pleurodema*. *Ann. Mag. Nat. Hist.*, 9 (20): 450-478.
- PARKER, H. W. (1927): The brevicipitid frogs allied to the genus *Gastrophryne*. *Occ. Pap. Mus. Zool. Univ. Michigan*, 187: 1-6.
- PARKER, H. W. (1928): Reptiles and batrachians from Mato Grosso and E. Bolivia. *Ann. Mag. Nat. Hist.*, 10 (2): 96-99.
- PARKER, H. W. (1934): *A monograph of frogs of the family Microhylidae*. British Museum (Natural History), London, 208 pp.
- PARKER, H. W. (1940): Lake Titicaca amphibia (The Percy Sladen Trust Expedition). *Trans. Linnean Soc. London*, 1 (2): 202-216.
- PÉFAUR, J. E. & DUELLMAN, W. E. (1980): Community structure in high Andean herpetofaunas. *Trans. Kansas Acad. Sci.*, 83: 45-65.
- PERACCA, M. G. (1897): Viaggio del Dott. Alfredo Borelli nel Chaco boliviano e nella Repubblica Argentina. *Boll. Mus. Zool. Anat. comp. R. Univ. Torino*, 12 (274): 1-19.
- PÉREZ, M. E. (1997): Una evaluación preliminar de los anfibios y reptiles de las Pampas del Heath (Provincia Iturralde, Departamento La Paz). *Ecología en Bolivia*, 30: 43-54.
- PÉREZ, M. E. (1998): *Dieta y ciclo gametogénico anual de Telmatobius culeus (Anura: Leptodactylidae) en el Lago Titicaca (Huiñaimarca)*. MS. Thesis, Universidad Mayor de San Andrés, La Paz. 150 pp. (unpublished).
- PROCTER, W. B. (1921): On a small collection of reptiles and batrachians made by Mr. Goodfellow in eastern Bolivia (1918-1919). *Ann. Mag. Nat. Hist.*, 9 (7): 189-192.
- REBOLLEDO, P. & TAPIA, M. C. (1992): *La herpetofauna de la Reserva de Vida Silvestre de los Ríos Blanco y Negro: diversidad, distribución, pautas para su conservación y uso sostenible*. 30 pp. Museo de Historia Natural "Noel Kempff Mercado", Santa Cruz (unpublished).
- REBOLLEDO, P. & MONTAÑO, M. E. (coord.) (1996): *Primera evaluación de la ictio y herpetofauna de la Laguna Cáceres y áreas de influencia en el Pantanal Boliviano (Germán Busch-Santa Cruz)*. Museo de Historia Natural "Noel Kempff Mercado", Santa Cruz (unpublished).
- REICHLE, S. (1996): *Freilandökologische Untersuchungen zur Amphibienfauna einer Savannenabflußrinne (Curichi) in Bolivien unter besonderer Berücksichtigung der Bioakustik*. Diplomarbeit, Univ. Hohenheim. 97 pp. (unpublished).
- REICHLE, S. (1997a): Frösche des Savannengebietes der Estación Biológica del Beni (EBB), Bolivien. Teil I: Einleitung, Untersuchungsgebiet, Artenliste der Anuren des Schutzgebietes und die Familien Bufonidae und Pseudidae. *Herpetofauna*, 19 (106): 5-11.
- REICHLE, S. (1997b): Frösche des Savannengebietes der Estación Biológica del Beni (EBB), Bolivien. Teil III: Die Familie Hylidae (Teil 2), Schlußbetrachtung und Literatur. *Herpetofauna*, 19 (108): 11-18.
- REICHLE, S. (1997c): Frösche des Savannengebietes der Estación Biológica del Beni (EBB), Bolivien. Teil II: Die Familien Leptodactylidae, Microhylidae und Hylidae (Teil 1). *Herpetofauna*, 19 (107): 11-18.
- REICHLE, S. (1999): Zur Kenntnis dreier Leptodactyliden aus Bolivien. *Ischnocnema sanctaecrucis* Harvey & Keck, 1995,

- Eleutherodactylus fenestratus* (Steindachner, 1864) und *Eleutherodactylus samaipatae* Köhler & Jungfer, 1995. *Herpetofauna*, 21 (123): 5-9.
- REICHLE, S., GUERRA, F. & PÉREZ, E. (1997): Geographic distribution: *Phyllomedusa bicolor*. *Herp. Rev.*, 28 (1): 48.
- REICHLE, S. & KÖHLER, J. (1996a): Erstnachweis von drei Froscharten für Bolivien: *Hyla tritaeniata*, *Phyllomedusa palliata* und *Colostethus marchesianus*. *Herpetofauna*, 18 (101): 32-34.
- REICHLE, S. & KÖHLER, J. (1996b): Geographic distribution: *Caecilia marcusi*. *Herp. Rev.*, 27 (4): 208.
- REICHLE, S. & KÖHLER, J. (1997): A new species of *Eleutherodactylus* (Anura: Leptodactylidae) from the Andean slopes of Bolivia. *Amphibia-Reptilia*, 18: 333-337.
- REICHLE, S. & KÖHLER, J. (1998): Saisonale und wasserstandsabhängige Rufplatzverteilung von Froschlurchen der südlichen Beni-Savannen, Bolivien. *Salamandra*, 34 (1): 43-54.
- REICHLE, S. & LÖTTERS, S. (2000): Amphibiensforschung in Bolivien: Von Hundert auf Zweihundert in Zehn.- DATZ Sonderheft "Frösche und Kröten", Stuttgart, pp. 16-19.
- REICHLE, S., LÖTTERS, S. & DE LA RIVA, I. (2001): A new species of the *discoidalis* group of *Eleutherodactylus* (Anura, Leptodactylidae) from inner-Andean valleys of Bolivia, with notes on the variation of the advertisement call of *Eleutherodactylus cruralis*. *J. Herpetol.* (in press).
- REICHLE, S., LÖTTERS, S. & MALDONADO, M. (2000): Two records of salamanders, genus *Bolitoglossa*, from Bolivia, providing southern range extension for the order (Amphibia: Caudata: Plethodontidae). *Herp. Nat. Hist.* (in press).
- REYNOLDS, R. P. & FOSTER, M. S. (1992): Four new species of frogs and one new species of snake from the Chapare region of Bolivia, with notes on other species. *Herpetol. Monogr.*, 6: 83-104.
- RIVERO, J. A. (1961): Salientia of Venezuela. *Bull. Mus. Comp. Zool.*, 126: 1-207.
- RIVERO, J. A. (1988): Sobre las relaciones de las especies del género *Colostethus* (Amphibia, Dendrobatidae). *Mem. Soc. Cienc. Nat. La Salle*, 48 (129): 2-32.
- RODRÍGUEZ, L. O. & MYERS, C. W. (1993): A new poison frog from Manu National Park, Peru (Dendrobatidae, *Epipedobates*). *Amer. Mus. Novit.*, 3068: 1-15.
- SANCHÍZ, B. & DE LA RIVA, I. (1993): Remarks on the tarsus of centrolenid frogs (Amphibia, Anura). *Graellsia*, 49: 115-117.
- SARMIENTO, J., BARRERA, S., BERNAL, N. & APARICIO, J. (1996): Fauna de una localidad del Altiplano central Huajara, Departamento de Oruro (Bolivia). *Rev. Bol. de Ecol.* 1 (1): 21-28.
- SAVAGE, J. M. (1969): Clarification of the status of the toad *Bufo veraguensis* O. Schmidt, 1857. *Copeia*, 1969 (1): 178-179.
- SAVAGE, J. M. (1972): The systematic status of *Bufo simus* O. Schmidt, with description of a new toad from western Panama. *J. Herpetol.*, 6 (1): 25-33.
- SAVAGE, J. M. & CEI, J. M. (1965): A review of the leptodactylid frog genus, *Odonophryne*. *Herpetologica*, 21 (3): 178-195.
- SAVAGE, J. M. & HEYER, W. R. (1969): The tree frogs (Family Hylidae) of Costa Rica: diagnosis and distribution. *Rev. Biol. Trop.*, 16 (1): 1-127.
- SCHMIDT, K. P. (1954): Notes on frogs of the genus *Telmatobius* with description of two new Peruvian species. *Fieldiana Zool.*, 34: 277-287.
- SCHMIDT, K. P. & INGER, R. F. (1951): Amphibians and reptiles of the Hopkins-Branner expedition to Brazil. *Fieldiana Zool.*, 31 (42): 439-465.
- SCHMIDT, O. (1857): Diagnosen neuer Frösche des zoologischen Cabinets zu Krakau. *Math. Natwiss. Cl. Kaiserl. Akad. Wiss. Wien*, 24: 10-15.
- SCHMIDT, O. (1858): Deliciae Herpetologicae Musei Zoologici Cracoviensis. Beschreibung

- der im K. K. Museum zu Krakau befindlichen und von L. v. Warszewicz in Neu-Granada und Bolivia gesammelten ungeschwänzten Batrachier. *Denkschr. Math.-Naturwiss. Cl. Kaiserl. Akad. Wien*, 14: 237-258.
- SCROCCHI, G. J. (1995): *Informe sobre la herpetofauna del Parque Nacional Noel Kempff Mercado, Departamento Santa Cruz, Bolivia*. FAN-TNC (unpublished).
- SHREVE, B. (1959): A new *Phyllomedusa* from Bolivia (Salientia, Hylidae). *Breviora*, 113: 114.
- SILVERSTONE, P. A. (1976): A revision of the poison-arrow frogs of the genus *Phyllobates* Bibron in Sagra (family Dendrobatiidae). *Nat. Hist. Mus. Los Angeles Co. Sci. Bull.*, 27: 1-53.
- SINSCH, U., SALAS, A. W. & CANALES, V. (1995): Reassessment of central Peruvian Telmatobiinae (genera *Batrachophryalus* and *Telmatobius*). I. Morphometry and classification. *Alytes*, 13 (1): 14-44.
- SMITH, E. N., REICHLE, S. & HARVEY, M. B. (2000). A new species of *Osteocephalus* (Anura, Hylidae) from Bolivia, with comments on the taxonomic status of *O. ayarzaguai* Gorzula & Señaris, 1996 and *O. yasuni* Ron & Pramuk, 1999. *Alytes* (in press).
- STEINDACHNER, F. (1892): Über zwei noch unbeschriebene *Nototrema*-Arten aus Ecuador und Bolivia. *Sitzber. Akad. Wiss. Wien*, 100: 337-342.
- SUÁREZ, S. & MURILLO, C. (1990): *Estudios básicos sobre flora y fauna silvestre del Parque Nacional Amboró. Lista de reptiles y anfibios del área del Parque Amboró*. Plan Maestro Parque Nacional Amboró, MACABID, Buenavista, Santa Cruz, 5 pp. (unpublished).
- TABER, A., NAVARRO, G. & ARRIBAS, M. A. (1997): A new park in the Bolivian Gran Chaco - an advance in tropical dry forest conservation and community-based management. *Oryx*, 13 (3): 189-198.
- TAYLOR, E. H. (1968): *The caecilians of the world. A taxonomic review*. Univ. Kansas Press. 848 pp.
- TRAVASSOS, L., TRAVASSOS, H., REGO-BARRROS, A., ALBURQUERQUE, D., OLIVEIRA, S., LEMOS, A. & LOPER, H. (1957): Exursão científica realizada nas zonas das Estradas de Ferro Noroeste do Brasil e Brasil-Bolivia em janeiro e fevereiro de 1955. *Publ. Avul. Mus. Nac.*, 20: 1-44.
- TRUEB, L. (1974): Systematic relationships of Neotropical horned frogs, genus *Hemiphractus* (Anura: Hylidae). *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 29: 1-60.
- TRUEB, L. & CANNATELLA, D. C. (1986): Systematics, morphology, and phylogeny of genus *Pipa* (Anura: Pipidae). *Herpetologica*, 42 (4): 412-449.
- TRUEB, L. & DUELLMAN, W. E. (1971): A synopsis of the Neotropical hylid frogs, genus *Osteocephalus*. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas*, 1: 1-47.
- TSCHUDI, J. J. VON (1838): Classification der Batrachier, mit Berücksichtigung der fossilen Thiere dieser Abteilung der Reptilien. *Mem. Soc. Sci. Nat. Neuchâtel*, 2: 1-99.
- VACA, N. (1993): *Informe del estudio herpetológico de la Reserva de Vida Silvestre de los Ríos Blanco y Negro*. Informe trimestral de alcance. Proyecto Prioridades de Conservación, 11 pp. (unpublished).
- VACA, N. (1994): *Relevamiento de la flora y fauna del territorio indígena Parque Nacional Isiboro Sécure (1993)*. Herpetología. TIPNIS, CIDDEBENI, FIA: 65-66. (Unpublished).
- VELLARD, J. (1951): Estudios sobre batracios andinos. I. El grupo *Telmatobius* y formas afines. *Mem. Mus. Hist. Nat. "J. Prado"*, 1: 3-89.
- VELLARD, J. (1953): Estudios sobre batracios andinos. II.- El grupo *Marmoratus* y formas afines. *Mem. Mus. Hist. Nat. "J. Prado"*, 2: 1-53.
- VELLARD, J. (1955): Répartition des batraciens dans les Andes au sud de l'équateur. *Trav. Inst. Franç. Ét. Andines*, 5: 141-161.

- VELLARD, J. (1957): Estudios sobre batracios andinos. IV. El género *Gastrotheca*. *Mem. Mus. Hist. Nat. "J. Prado,"* 5: 1-47.
- VELLARD, J. (1960a): Estudios sobre batracios andinos. VI.- Notas complementarias sobre *Telmatobius*. *Mem. Mus. Hist. Nat. "J. Prado,"* 10: 1-19.
- VELLARD, J. (1960): Estudios sobre batracios andinos. VII. El género *Pleurodema* en los Andes Peruanos. *Mem. Mus. Hist. Nat. "J. Prado,"* 10 (7): 1-14.
- VELLARD, J. (1969): Les *Telmatobius* du group *marmoratus*. *Bull. Mus. Natn. Hist. Nat.,* 40 (6): 1110-1113.
- VELLARD, J. (1970): Contribución al estudio de los batracios andinos. *Rev. Mus. Arg. Cien. Nat. "B. Rivadavia,"* 10 (1): 1-21.
- VELLARD, J. (1992): Associated animal communities. The amphibia, pp. 449-457, in: Dejoux, C. & Iltis, A. (eds.), *Lake Titikaka. A synthesis of limnological knowledge*, Kluwer Academic Publishers.
- VELOSO, A. & NAVARRO, J. (1988): Lista sistemática y distribución geográfica de anfibios y reptiles de Chile. *Boll. Mus. reg. Sci. nat. Torino,* 6 (2): 481-539.
- WAKE, D. B., BRAME, A. H. & THOMAS, R. (1982): A remarkable new species of salamander allied to *Bolitoglossa altamazonica* (Plethodontidae) from southern Peru. *Occ. Pap. Mus. Zool. Louisiana State Univ.,* 58: 1-21.
- WAKE, M. H. (1984): A new species of *Caecilia* (Amphibia: Gymnophiona) from Bolivia. *Amphibia-Reptilia,* 5: 215-220.
- WALKER, C. F. (1973): A new genus and species of microhylid frog from Ecuador. *Occ. Pap. Mus. Nat. Hist. Univ. Kansas,* 20: 1-7.
- WERNER, F. (1899): Beschreibung neuer Reptilien und Batrachier. *Zool. Anz.,* 22: 479-484.
- WERNER, F. (1901): Wissenschaftliche Mitteilungen. 1. Beschreibung neuer Frosche aus Bolivia, Ostindien und Neu Guinea. *Zool. Anz.,* 24: 97-103.

Recibido: 28/09/99

Aceptado: 13/01/00

Color Photographs.— Abbreviations of authors of the photographs are as follows: IR, I. De la Riva; JD, J. Dorda; WED, W. E. Duellman (courtesy of the Natural History Museum, The University of Kansas); MG, M. Göd; PI, P. Ibisch; KHJ, K.-H. Jungfer; JK, J. Köhler; AK, A. Kwet; JL, J. Lynch (courtesy of the Natural History Museum, The University of Kansas); AM, A. Mújica; SR, S. Reichle; and EW, E. Wild.

Fotografías en color.— Las abreviaturas de los autores de las fotografías son: IR, I. De la Riva; JD, J. Dorda; WED, W. E. Duellman (courtesy of the Natural History Museum, The University of Kansas); MG, M. Göd; PI, P. Ibisch; KHJ, K.-H. Jungfer; JK, J. Köhler; AK, A. Kwet; JL, J. Lynch (courtesy of the Natural History Museum, The University of Kansas); AM, A. Mújica; SR, S. Reichle; y EW, E. Wild



Atelopus tricolor. Chapare, Cochabamba (JK)



Bufo amboroensis. Río Chua Khocha, Cochabamba (IR)



Bufo arenarum. Vaca Guzmán, Chuquisaca (JK)



Bufo castaneoticus. Cobija, Pando (JK)



Bufo granulosus. Rurrenabaque, Beni (SR)



Bufo guttatus. P. N. Noel Kempff Mercado, Santa Cruz (SR)



Bufo justinianoi. Chapare, Cochabamba (JK)



Bufo "margaritifer". La Bola, Santa Cruz (IR)



Bufo marinus. Valle de Sajta, Cochabamba (IR)



Bufo paracnemis. Puerto Almacén, Santa Cruz (IR)



Bufo poeppigii. Carahuasi, Cochabamba (IR)



Bufo quechua. Incachaca, Cochabamba (JK)



Bufo spinulosus. López Mendoza, Cochabamba (IR)



Bufo veraguensis. La Yunga, Santa Cruz (IR)



Dendrophryniscus minutus. Valle de Sajta, Cochabamba (IR)



Melanophryniscus rubriventris. López Mendoza, Cochabamba (IR)



Cochranella bejaranoi. Sehuencas, Cochabamba (JK)



Cochranella nola. El Fuerte de Samaipata, Santa Cruz (JK)



Cochranella pluvialis. Coroico, La Paz (SR)




Hyalinobatrachium bergeri. Chapare, Cochabamba (JK)



Allobates femoralis. Chalalán, La Paz, (SR)



Colostethus brunneus. P. N. Noel Kempff, Santa Cruz (SR)



Colostethus mcdiarmidi. Río Yuyo, La Paz (IR)



Colostethus trilineatus. Estación Biológica Beni, Beni (SR)



Epipedobates bolivianus. Correo, La Paz (IR)



Epipedobates hahneli. Cobija, Pando (JK)



Epipedobates pictus. P. N. Noel Kempff Mercado, Santa Cruz (IR)



Epipedobates trivittatus. San Sebastián, Pando (SR)



Gastrotheca lauzuricae. Serranía Siberia, Cochabamba (IR)



Gastrotheca marsupiata. Sehuencas, Cochabamba (JK)



Gastrotheca splendens. Abra de la Cruz, Santa Cruz (IR)



Gastrotheca testudinea. Chapare, Cochabamba (JK)



Hyla acreana. Mataracú, Santa Cruz (SR)



Hyla albonigra. Toro Toro, Potosí (PI)



Hyla albopunctata. P. N. Noel Kempff Mercado, Santa Cruz (IR)



Hyla andina. Aguirre, Cochabamba (IR)



Hyla armata. Chaco, La Paz (IR)



Hyla balzani. Coroico, La Paz (SR)



Hyla bifurca. Puerto Almacén, Santa Cruz (IR)



Hyla boans. P. N. Noel Kempff Mercado, Santa Cruz (SR)



Hyla calcarata. San Sebastián, Pando (SR)



Hyla charazani. Charazani, La Paz (IR)



Hyla chlorostea (holotype). Paracti, Cochabamba (IR)



Hyla fasciata. Cobija, Pando (JK)



Hyla geographica. Puerto Almacén, Santa Cruz (IR)



Hyla granosa. Cobija, Pando (JK)



Hyla koechlini. Rurrenabaque, Beni (SR)



Hyla lanciformis. Valle de Sajta, Cochabamba (IR)



Hyla leali. Estación Biológica Beni, Beni (SR)



Hyla leucophyllata. Estación Biológica Beni, Beni (SR)



Hyla marianitae. Carahuasi, Cochabamba (IR)



Hyla marmorata. "Inselberg" Santa Cruz (JK)



Hyla melanargyrea. P. N. Noel Kempff Mercado, Santa Cruz (SR)



Hyla minuta. Laguna Volcán, Santa Cruz (SR)



Hyla nana. Estación Biológica Beni, Beni (SR)



Hyla parviceps. Puerto Almacén, Santa Cruz (IR)



Hyla punctata. Puerto Almacén, Santa Cruz (IR)



Hyla raniceps. La Bola, Santa Cruz (IR)




Hyla rhodopepla. Mataracú, Santa Cruz (SR)



Hyla riveroi. Cobija, Pando (JK)



Hyla sarayacuensis San Sebastián, Pando (SR)



Hyla schubarti. Cuzco Amazónico, Peru (EW)



Hyla tritaeniata. Estación Biológica Beni, Beni (SR)



Osteocephalus buckleyi. Mataracú, Santa Cruz (SR)



Osteocephalus cf. leprieurii. Los Fierros, Santa Cruz (SR)



Osteocephalus pearsoni. Rurrenabaque, Beni (SR)



Osteocephalus taurinus. Estación Biológica Beni, Beni (SR)



Phrynohyas coriacea. Puerto Almacén, Santa Cruz (IR)



Phrynohyas resinifictrix. Centro Científico Yasuní, Napo, Ecuador (KHJ)



Phrynohyas venulosa. P. N. Noel Kempff Mercado, Santa Cruz (IR)




Scarthyla ostinodactyla. Iquitos, Peru (MG)



Scinax acuminatus. Lago Ipacaraí, Paraguay (KHJ)



Scinax castroviejoi. Laguna Volcán, Santa Cruz (IR)



Scinax chiquitanus. Puerto Almacén, Santa Cruz (AM)



Scinax fuscovarius. Puerto Almacén, Santa Cruz (IR)



Scinax garbei. Cobija, Pando (JK)



Scinax nasicus. Estación Biológica Beni, Beni (SR)



Scinax nebulosus. P. N. Noel Kempff Mercado, Santa Cruz (SR)



Scinax parkeri. P. N. Noel Kempff Mercado, Santa Cruz (IR)



Scinax ruber. Estación Biológica Beni, Beni (SR)



Scinax squalirostris. Apolo, La Paz (IR)



Sphaenorhynchus lacteus. Puerto Almacén, Santa Cruz (IR)



Phyllomedusa bicolor. Pingo de Oro, Pando (SR)



Phyllomedusa boliviiana. P. N. Amboró, Santa Cruz (IR)



Phyllomedusa camba. Puerto Almacén, Santa Cruz (IR)



Phyllomedusa hypochondrialis. Buenavista, Santa Cruz (IR)



Phyllomedusa palliata. Estación Biológica Beni, Beni (SR)



Phyllomedusa sauvagii. Raco, Tucumán, Argentina (IR)



Phyllomedusa tomopterna. Cuzco Amazónico, Perú (IR)



Phyllomedusa vaillanti. Mataracú, Santa Cruz (JK)



Ceratophrys cornuta. Puerto Almacén, Santa Cruz (IR)



Ceratophrys cranwelli. Santa Cruz de la Sierra, Santa Cruz (JK)



Chacophrys pierottii. Captive specimen from the pet trade (IR)



Lepidobatrachus laevis. Cerro Cortado, Santa Cruz (SR)



Adenomera andreae. Cobija, Pando (JK)



Adenomera diptyx. Santa Cruz de la Sierra, Santa Cruz (JK)



Adenomera hylaedactyla. Cobija, Pando (JK)



Leptodactylus boliviensis. Puerto Almacén, Santa Cruz (IR)



Leptodactylus bufonius. La Bola, Santa Cruz (IR)



Leptodactylus "chacensis-macrosternum". Vaca Guzmán, Chuquisaca (JK)



Leptodactylus didymus. Cobija, Pando (JK)



Leptodactylus elenae. Macuñucu, Santa Cruz (JK)



Leptodactylus fuscus. Buenavista, Santa Cruz (JK)



Leptodactylus gracilis. El Fuerte de Samaipata, Santa Cruz (JK)



Leptodactylus griseigularis. Chapare, Cochabamba (JK)



Leptodactylus knudseni. Rurrenabaque, Beni (SR)



Leptodactylus labyrinthicus. Puerto Almacén, Santa Cruz (IR)



Leptodactylus laticeps. Argentina (WED)



Leptodactylus latinasus. Arroyo Los Naranjos, Jujuy, Argentina (WED)



Leptodactylus leptodactyloides. P. N. Noel Kempff Mercado, Santa Cruz (IR)



Leptodactylus mystaceus. Río Chevejecure, Beni (SR)



Leptodactylus mystacinus. Paraguay (JK)



Leptodactylus ocellatus. Lago Ipacaraí, Paraguay (KHJ)



Leptodactylus pentadactylus. Isla Coiba, Panama (IR)



Leptodactylus petersi. Buenavista, Santa Cruz (JK)



Leptodactylus podicipinus. San Ramón, Santa Cruz (JK)



Leptodactylus rhodomystax. Manaus, Brazil (KHJ)



Leptodactylus rhodonotus. La Hoyada, Santa Cruz (JK)



Leptodactylus syphax. "Inselberg", Santa Cruz (JK)



Lithodytes lineatus. Mataracú, Santa Cruz (SR)



Physalaemus albonotatus. Santa Cruz de la Sierra, Santa Cruz (JK)



Physalaemus biligonigerus. Santa Cruz de la Sierra, Santa Cruz (IR)



Physalaemus cuvieri. Porto Alegre, Brazil (AK)



Physalaemus nattereri. Santa Rosa de la Roca, Santa Cruz (JK)



Physalaemus petersi. Mataracú, Santa Cruz (JK)




Pleurodema cinereum. Huayllamarca, Oruro (IR)



Pleurodema guayapae. Izozog, Santa Cruz (SR)



Pleurodema marmoratum. Comanche, La Paz (IR)



Pseudopaludicola boliviensis. Estación Biológica Beni, Beni (SR)



Pseudopaludicola mystacalis. P. N. Noel Kempff Mercado, Santa Cruz (IR)



Vanzolinia discodactylus. Cobija, Pando (JK)



Eleutherodactylus ashkapara. Chapare, 2100 m, Cochabamba (JK)



Eleutherodactylus cruralis. Mataracú, Santa Cruz (SR)



Eleutherodactylus danae. Chapare, Cochabamba (JK)



Eleutherodactylus discoidalis. La Yunga, Santa Cruz (IR)



Eleutherodactylus fenestratus. Santa Cruz del Valle Ameno, La Paz (IR)



Eleutherodactylus fraudator. Serranía Siberia, Cochabamba (IR)



Eleutherodactylus llojsintuta. Sehuencas, Cochabamba (JK)



Eleutherodactylus mercedesae. Road Caranavi-Yucumo, La Paz (SR)



Eleutherodactylus olivaceus. Chapare, Cochabamba (JK)



Eleutherodactylus peruvianus. Santa Cruz del Valle Ameno, La Paz (IR)



Eleutherodactylus platydactylus. Sehuencas, Cochabamba (JK)



Eleutherodactylus pluvicanorus. Carahuasi, Cochabamba (SR)



Eleutherodactylus rhabdolaemus. Río Ronquito, Cochabamba (JK)



Eleutherodactylus samaipatae. El Fuerte de Samaipata, Santa Cruz (JK)



Eleutherodactylus toftae. Paractito, Cochabamba (JD)



Eleutherodactylus ventrimarmoratus. Mera, Ecuador (JL)



Eleutherodactylus zongoensis. Zongo Valley, La Paz (SR)



Ischnocnema quixensis. P. N. Amacayacu, Colombia (IR)



Ischnocnema sanctaerucis. Carahuasi, Cochabamba (JK)



Odontophrynus americanus. Pampa Grande, Santa Cruz (SR)



Odontophrynus lavillai. Santa Cruz de la Sierra, Santa Cruz (JK)



Phrynobatrachus kempffii. Serranía Siberia, Cochabamba (IR)



Phrynobatrachus laplacai. Unduavi, La Paz (SR)



Phrynobatrachus pinguis (paratypes). Choquetanga Chico, La Paz (IR)



Phyllonastes carrascoicola. Chapare, Cochabamba (JK)



Phyllonastes ritarasquinae. Chapare, 1250 m, Cochabamba (JK)



Telmatobius boliviensis. Chaco, La Paz (IR)



Telmatobius culeus. Lago Titicaca, La Paz (KHJ)



Telmatobius edaphonastes. Serranía Siberia, Santa Cruz (SR)



Telmatobius huayra (holotype). Campamento Khastor, Potosí (IR)



Telmatobius jahuir. Kkota Pata, La Paz (IR)



Telmatobius marmoratus. Comanche, La Paz (IR)



Telmatobius simonsi. Sucre, Chuquisaca (JK)



Telmatobius verrucosus (holotype). Chaco, La Paz (IR)



Telmatobius yuracare. Sehuencas, Cochabamba (JD)



Chiasmocleis albopunctata. Río Seco, Santa Cruz (JK)



Chiasmocleis ventrimaculata. Estación Biológica Beni, Beni (SR)



Dermatonotus muelleri. Paraguay (JK)



Elachistocleis bicolor. Estación Biológica Beni, Beni (SR)



Elachistocleis ovalis. Río Seco, Santa Cruz (JK)



Hamptophryne boliviiana. Puerto Almacén, Santa Cruz (JD)



Pipa pipa. Captive specimen from the pet trade (IR)



Lysapsus limellus. P. N. Noel Kempff, Santa Cruz (JK)



Pseudis paradoxa. Montero, Santa Cruz (JK)



Rana palmipes. Paracti, Cochabamba (JK)



Bolitoglossa altamazonica. Villa Tunari, Cochabamba (SR)



Caecilia marcusii. Totaizal, Beni (SR)



Siphonops annulatus. Puerto Maldonado, Peru (WED)



Siphonops paulensis. Santa Cruz de la Sierra, Santa Cruz (JK)